

Magnetic Vortex Dynamics

in a 2D easy plane ferromagnet

by

Lara Thompson

B.Sc., The University of Waterloo, 2002

B.Math., The University of Waterloo, 2001

A THESIS SUBMITTED IN PARTIAL FULFILMENT OF
THE REQUIREMENTS FOR THE DEGREE OF

MASTER OF SCIENCE

in

THE FACULTY OF GRADUATE STUDIES

(Department of Physics and Astronomy)

We accept this thesis as conforming
to the required standard

.....
.....

THE UNIVERSITY OF BRITISH COLUMBIA

February 2, 2008

© Lara Thompson, 2004

Abstract

In this thesis, we consider the dynamics of vortices in the easy plane insulating ferromagnet in two dimensions. In addition to the quasiparticle excitations, here spin waves or magnons, this magnetic system admits a family of vortex solutions carrying two topological invariants, the winding number or vorticity, and the polarization.

A vortex is approximately described as a particle moving about the system, endowed with an effective mass and acted upon by a variety of forces. Classically, the vortex has an inter-vortex potential energy giving a Coulomb-like force (attractive or repulsive depending on the relative vortex vorticity), and a gyrotropic force, behaving as a self-induced Lorentz force, whose direction depends on both topological indices.

Expanding semiclassically about a many-vortex solution, the vortices are quantized by considering the scattered magnon states, giving a zero point energy correction and a many-vortex mass tensor. The vortices cannot be described as independent particles—that is, there are off-diagonal mass terms, such as $\frac{1}{2}M_{ij}v_i v_j$, that are non-negligible.

This thesis examines the full vortex dynamics in further detail by evaluating the Feynman-Vernon influence functional, which describes the evolution of the vortex density matrix after the magnon modes have been traced out. In addition to the set of forces already known, we find new damping forces acting both longitudinally and transversely to the vortex motion. The vortex motion within a collective cannot be entirely separated: there are damping forces acting on one vortex due to the motion of another. The effective damping forces have memory effects: they depend not only on the current motion of the vortex collection but also on the motion history.

Contents

Abstract	iii
Contents	v
List of Figures	vii
Acknowledgements	ix
1 Introduction	1
1.1 Symmetry breaking	2
1.2 Classical Solitons	5
1.3 Quantum Solitons	7
1.3.1 The particle theorists	7
1.3.2 In condensed matter theory	9
1.3.3 Superfluid He^4	12
1.3.4 Magnetic vortices	15
1.4 Easy plane insulating ferromagnet	17
2 Magnons	21
2.1 Magnon equations of motion	21
2.2 Quantum propagator	24
2.2.1 Spectrum via tracing over the propagator	26
2.3 Thermal equilibrium density matrix	27
2.3.1 Magnon density matrix	27
2.4 Summary	29
3 Vortices	31
3.1 Force between vortices	33
3.2 The gyrotropic force and the vortex momentum	36
3.2.1 The gyrotropic force	36
3.2.2 The vortex momentum	39
3.3 Motion of vortex pairs	42
3.4 Vortex mass	43
3.5 Quantization of magnetic vortices	44
3.5.1 Phase shifts in the Born approximation	46
3.5.2 Bound modes	50

4 Vortex dynamics	53
4.1 Vortex-magnon interaction terms	54
4.2 Perturbation theory results	55
4.2.1 Vortex mass revisited	55
4.2.2 Radiation of magnons	59
4.2.3 Zero point energy	63
4.3 Vortex influence functional	64
4.3.1 Quantum Brownian motion	67
4.3.2 Semiclassical solution of perturbed magnons	69
4.3.3 Evaluating the influence functional	71
4.3.4 Interpreting the imaginary part	73
4.3.5 Interpreting the real part	76
4.4 Discussion of vortex effective dynamics	78
4.4.1 Comparison with radiative dissipation	79
4.4.2 Extending results to many vortices	79
4.4.3 Frequency dependent motion	82
4.4.4 Summary	83
5 Conclusions	85
5.1 Open questions	86
A Some mechanics	87
A.1 Imaginary time path integral	89
B Quantization of classical solutions	91
B.1 Quantizing soliton solutions	92
B.1.1 In a path integral formalism	94
B.1.2 Collective coordinates	97
C Spin path integrals	99
C.1 The semiclassical approximation	100
C.1.1 Coherent state path integral	101
C.1.2 Spectrum of a ferromagnetic plane of spins	104
Bibliography	107

List of Figures

- 1.1 Left, an example potential of a 1D field ϕ with a doubly degenerate ground state; right, an example potential. 5
- 1.2 A 2D XY-ferromagnet with a vortex connecting the degeneracy of spin directions. The central red dot signifies the vortex core. 5
- 1.3 A magnetic vortex formed by Heisenberg spins can be continuously deformed away by expanding about a point. 5
- 1.4 A vortex with $+1$ winding in a 2D Heisenberg ferromagnet with spins lying preferentially in the plane. 5
- 1.5 Re-enactment of the 1834 ‘first’ soliton sighting on the Union Canal near Edinburgh by John Scott Russell. 5
- 1.6 An illustrative potential of a one dimensional particle. A soliton is analogous to the second minimum at $x = 0$ 5
- 1.7 The two degenerate dimer states of *trans*-(CH) $_N$, polyacetylene. 10
- 1.8 The band structure of polyacetylene, gapped due to the electron-phonon interactions. Note the two isolated bands. 10
- 1.9 A kink solution connecting the two degenerate dimer ground states, shown, left, on the linear polyacetylene chain. 10
- 1.10 The equi-pressure lines of a fluid surrounding a rotating cylinder. The pressure differential top and bottom. 10

- 2.1 A comparison of the easy plane magnon spectrum and density of states with the regular isotropic ferromagnet. 10

- 3.1 Vortex spin configuration: left, a vortex with $q = -1$; right, a vortex with $q = 1$ 33
- 3.2 Two vortex spin configurations. Left, two vortices with $q = 1$; right, vortices with $q = 1$ and $q = -1$; both vortices with $q = 1$ 33
- 3.3 Intervortex forces: top, two vortices of opposite vorticity attract; bottom, two vortices with same sense vorticity repel. 33
- 3.4 The spin path mapped onto the unit sphere. The area traced out by its motion gives the Berry’s phase. 37
- 3.5 The gyrotropic force: left, a vortex with $p = 1$ and $q = -1$ traveling to the right experiences an upward force; right, a vortex with $p = 1$ and $q = 1$ traveling to the right experiences a downward force. 37
- 3.6 Sequenced photographs of a pair of fluid vortices with same sense vorticity. Photos were taken at 2 second intervals. 42
- 3.7 Sequenced photographs of a pair of fluid vortices with opposite sense vorticity. Photos were taken at 4 minute intervals. 42
- 3.8 The directions relevant to a small translation of the vortex along $\delta\mathbf{r}$ 52

- 4.1 Lowest order contributing diagram for the first order vortex-magnon coupling term. 57
- 4.2 Definition of angles for evaluation of off-diagonal mass terms. 58
- 4.3 The dissipation rate from perturbation theory; first assuming infinite mass and then adding corrections due to finite mass. 58

- B.1 An illustrative potential of a one dimensional particle. 91

Acknowledgements

Thanks to Phil for choosing an excellent masters research topic. To my mom who read my thesis and corrected it despite not understanding every third word, although learning that equations have a grammar all their own! To Talie in Toronto for housing me in the midst of the crunch and showing me a good time otherwise to cool off. To Yan for sharing with me the mountains.

“What did the condensed matter theorist say to the soliton? As long as you aren’t empirical, you’re all right with me.” –Lateef Yang, August 11, 2004

Chapter 1

Introduction

In a wide variety of systems, there exist vortices, high energy states nonetheless significant in system dynamics at low temperatures. Despite its high energy, a vortex can nonetheless form via tunneling processes or at a boundary with only a small energy barrier. They are exceptionally stable, arguable topologically, and, in fact, can only be destroyed if one meets its ‘anti-vortex’ or, equivalently, annihilates at a boundary (where it has met its image vortex). Cooling a system down vortex-free is non-trivial, and, in general, we retain a low density of vortex states down to the lowest temperatures.

Quantum vortices were first proposed in the 1950’s in superfluid helium to explain the decay of persistent currents. Since then, they have been proposed and measured in, for example, superconductors and a variety of magnetic systems. The dynamics are well described phenomenologically as a point-like particle in 2D (or as a line in 3D) endowed with an effective mass and acted upon by a variety of forces. Microscopic derivations of the particle properties of a quantum vortex have been plagued by decades of debate and controversy. A recent resurgence in debates began in the 1990’s concerning the so-called Magnus force, a force borrowed from classical fluids acting perpendicular to the velocity. Ao and Thouless² claimed that in superfluid helium (He II) there is a universal form of this force, independent of quasiparticle scattering. Others argue that there should be, in addition to the bare Magnus force, a transverse damping force, reinforcing or opposing the Magnus force^{22,61,70}.

In this thesis, we consider a relatively simple magnetic system, a 2D insulating ferromagnet with easy plane anisotropy, admitting a family of topologically stable vortices. We derive microscopically the vortex effective mass and, in addition to the previously reported gyrotropic force, the magnetic analogue to the Magnus force, and inter-vortex Coulomb-like forces, we derive a variety of vortex damping forces. We find both the usual longitudinal damping force and a transverse damping that acts in combination with the gyrotropic force. A transverse damping force has not yet been considered in a magnetic system. In fact, all treatments of the dissipative motion of a vortex have been phenomenological, with the exception of Slonczewski’s⁵⁹ treatment with which we compare results in Chapter 4. A collection of vortices cannot be considered as a set of independent particles—they have mixed inertial terms and damping force terms.

We first review a few symmetry arguments for the existence and stability of vortex solutions. Besides revealing the similarity between vortices from various systems, we find that vortices are an example of a more general family of topological solitons.

We then briefly discuss the early work on quantizing solitons by the relativistic field theorists, focussing rather on the techniques than the various specific contributions. Note that we will use many of these techniques for quantizing the vortex in the easy plane magnetic system.

Next, we discuss briefly solitons in condensed matter systems and the exciting new phenomena found there. For example, by examining the conducting polymers, fractional charge was first predicted and observed.

Returning specifically to vortices, we briefly discuss the controversy in the microscopic derivation of the equations of motion for a superfluid vortex. This will introduce the variety of forces we should expect to act on a collection of vortices. Switching to magnetic systems, we find that despite the ease of direct experimental observation and simplicity of calculations not much work has been done here.

Finally, we introduce in detail the magnetic system under consideration. The symmetry of the system admits topologically stable vortices and gapless quasiparticles. The purpose of this thesis is to separate the quantum dynamics of the vortices from the effects of the perturbative quasiparticles, here magnons.

1.1 Symmetry breaking

Symmetry plays a crucial role in science and we strive to discover and exploit the symmetries of the laws of nature (Galilean or Lorentz invariance, gauge invariance, etc.). However, we find that the symmetry of physical states may be a smaller subset of the full symmetry in which it resides. For example, in a Heisenberg ferromagnet, we find a system of spins free to lie in any direction in 3D, preferring to align parallel to one another, however, in the absence of any magnetic fields, with no preference of which direction along which to lie. The ground state then chooses at random along what direction to align.

A system with a degenerate ground state is forced to spontaneously choose one state amid the degeneracy, an example of spontaneously broken symmetry. A discrete degeneracy is found in the problem of a field residing in a double well potential (as in Figure 1.1, left), or, more generally, an n -well potential. A continuous degeneracy in a system has a continuum of minima in the potential (as, for example, in Figure 1.1, right). The ferromagnet is an example of a system with a continuum of ground states, except that here, the potential is completely flat: there is no preference at all between directions.




Figure 1.1: Left, an example potential of a 1D field ϕ with a doubly degenerate ground state; right, an example potential of a 2D field with a continuum degeneracy in its ground state.

In general, different regions of a sample may choose different degenerate states or may even lie in an excited state. A mapping of the state taken across the sample, in all its available degrees of freedom, is called the order parameter. In a Heisenberg spin system, this is simply the spin vector in 3D as a function of position in the sample. The order parameter here can be mapped onto a unit sphere—a path along the sample is then traced as a path on the surface of the sphere. For a spin system confined to lie in the plane, the so-called XY model, the order parameter is mapped onto the unit circle.

Incidentally, the order parameter in superfluid helium II can also be mapped onto the unit circle so that it is topologically equivalent to the XY model. This does not mean, however, that the dynamics of the vortices in each system should be the same, but, rather, only that the topology of vortices is identical in the two systems.

If a system possesses discrete symmetries, to pass from one ground state to another there must be some transition region, or domain wall, separating different states. This domain wall, sometimes called a kink, is an example of a quasi-1D soliton.

For a continuous symmetry, we can imagine similar cases where certain regions are forced out of a ground state. As a simple example, consider the XY spin model. If the spins choose to nearly align along the boundary, turning very slowly so as to always radiate outward, as we near some central region the spins are less and less ferromagnetically aligned and, further, there is a point discontinuity at the very center (see Figure 1.2).

If we follow a path surrounding the vortex in order parameter space, that is along the unit circle, we find we must wrap around the unit circle once. This vortex is called a topological soliton with single *wrapping number* or *vorticity*. In this example, no matter how we smoothly deform the spins, we cannot continuously deform away this wrapping of the unit circle. We say that it is homotopically distinct from a zero winding path, or more simply a point.




Figure 1.2: A 2D XY-ferromagnet with a vortex connecting the degeneracy of spin directions. The central red dot signifies the point of discontinuity.




Figure 1.3: A magnetic vortex formed by Heisenberg spins can be continuously deformed away by expanding about a patch of the unit sphere not covered by the vortex path, shrinking the vortex to a point.

There are vortices with higher winding numbers, always integral to ensure continuity. Each family of solutions corresponding to a certain winding number is topologically stable. That is, there exists no homotopy, or continuous mapping, between solutions of differing winding numbers.

There are systems that admit vortices for which this topological stability is not guaranteed, and are thus not called topological solitons. Consider a general vortex residing on a sample for which the order parameter maps onto a unit sphere (Figure 1.3). The vortex is homotopically equivalent to a point (that is, a region with constant ground state) since we can imagine continuously shrinking the vortex away. In real space, this is equivalent to the ability of the spins to unwind, that is, all the spins twisting to all lie parallel to one another. Note that this unwinding is a special feature of the isotropy of the system. Although such a soliton does not possess topological stability, the entire plane must unwind, a macroscopic number of spins in the magnetic vortex case, so that the soliton is still essentially stable.

The vortices considered in this thesis have an order parameter lying on the unit sphere, however, with a higher potential at the north and south poles. They are very similar to the XY vortex shown in figure 1.2, except that the spins are not entirely restricted to lie in the plane and, at some energy expense to restore continuity, the spins twist out of plane at the vortex center choosing




Figure 1.4: A vortex with $+1$ winding in a 2D Heisenberg ferromagnet with spins lying preferentially in the plane.



Figure 1.5: Re-enactment of the 1834 ‘first’ soliton sighting on the Union Canal near Edinburgh by John Scott Russell.

spontaneously between the two possible perpendicular directions in which to twist. This direction is a second topological invariant of the vortices and is termed the *polarization*. An example of a vortex with unit winding number, or vorticity, and polarization out of the page is shown in Figure 1.4. There exist also zero polarization vortices lying entirely in the plane.

1.2 Classical Solitons

We found that vortices are examples of a topological solitons. Generally, a soliton is a finite energy localized solution of a wave equation, satisfying strict stability conditions under collisions with other soliton solutions[†].

[†]See, for instance, the excellent book by Rajaraman⁵⁰ on the quantization of solitons for a rigorous definition of a soliton.

The first reported soliton was in 1834 by John Scott Russell⁵³ in the Union Canal near Edinburgh (see Figure 1.5),

I was observing the motion of a boat which was rapidly drawn along a narrow channel by a pair of horses, when the boat suddenly stopped - not so the mass of water in the channel which it had put in motion; it accumulated round the prow of the vessel in a state of violent agitation, then suddenly leaving it behind, rolled forward with great velocity, assuming the form of a large solitary elevation, a rounded, smooth and well-defined heap of water, which continued its course along the channel apparently without change of form or diminution of speed. I followed it on horseback, and overtook it still rolling on at a rate of some eight or nine miles an hour, preserving its original figure some thirty feet long and a foot to a foot and a half in height. Its height gradually diminished, and after a chase of one or two miles I lost it in the windings of the channel. Such, in the month of August 1834, was my first chance interview with that singular and beautiful phenomenon which I have called the Wave of Translation.

He went on to build a 30' wave tank in his back garden in which to conduct further experiments on his "waves of translation".

In physics, there are the familiar optical solitons, with which demonstrations of long haul, low bit-error-rate transmissions have been made. In optics, a soliton is a localized EM wave with much higher power than a traditional optical signal. However, as opposed to regular low power optical transmissions, an optical soliton does not suffer dispersion, so that a signal is not distorted when transmitted over large distances.

A soliton is usually a solution to a partial differential equation in which competing non-linear terms cooperate to create a self-reinforcing large amplitude solution. For instance, for a non-linear dissipative system, ordinarily, wave solutions are dispersive, that is, different k modes separate, and dissipative, energy spreads in real space. For these special soliton solutions the two mechanisms can act in opposition so that the net result is a non-dispersive, non-dissipative wave.

More specifically, however, a vortex is an example of a topological soliton. These exist, not because of finely balanced non-linear terms in the equations of motion, but rather due to a degenerate freedom in the boundary conditions entailing the existence of homotopically distinct solutions (that is, solutions for which there is no continuous deformation from one to another).




Figure 1.6: An illustrative potential of a one dimensional particle. A soliton is analogous to the second minimum at $x = c$.

1.3 Quantum Solitons

1.3.1 The particle theorists

Solitons resemble extended particles, that is, they are non-dispersive localized packets of energy, even though they are solutions of non-linear wave equations. Elementary particles are localized packets of energy and are also believed to be solutions of some relativistic field theory. The particle theorists were thus highly motivated to find some quantum version of these classical solitons, that is, to quantize the solitons.

It isn't immediately clear how to make the correspondence between a classical soliton and some extended particle state of a quantized theory, or between any classical field solution and its quantum analogue for that matter. To understand the difficulty, consider first the simple case of a point particle in a potential. Classically, this particle has some definite position and momentum with some particular path chosen by its initial conditions. Quantum mechanically, the picture changes entirely! No longer can we associate a particle with a definite position and momentum; instead, we must describe the particle probabilistically via a wavefunction $\psi(x, t)$ giving the probability $|\psi(x, t)|^2$ to find the particle at point x and time t . How does one go from the soliton solution to some quantum wavefunction?

Procedures for establishing this correspondence developed in the mid-70's were essentially a generalization of the semiclassical expansion of non-relativistic quantum mechanics. It was shown that not only could we associate a quantum soliton-particle with the classical solution, but also a series of excited states by quantizing fluctuations about the soliton^{8,20}.

For a soliton, we quantize its motion by defining conjugate position $\mathbf{X}$ and momentum $\mathbf{P}$ operators and imposing commutation relations. In the original field, however, there is an entire continuum of degrees of freedom that remain.

These are taken up by the quasiparticle excitations.

The procedure is analogous to the quantization of a particle residing in a local minimum of the external potential (for example, $x = c$ in Figure 1.6). This local minimum is not the global minimum, and hence is not the true ground state; however, there is a potential barrier blocking it from decaying to the true ground state. This is the same for a soliton excitation, or a vortex, which is higher in energy than the ground state, however, stable against decay.

The quantization of the local minimum begins by assuming to zeroth order the classical solution, $x = c$. We expand the potential about this local minimum, finding quadratic behaviour to leading order, and proceed to quantize the perturbative excitations. Of course, a quadratic potential has simple harmonic excitations, so that the quantized solution can be envisioned as a hierarchy of simple harmonic excitations, centered, of course, about the classical minimum.

For a soliton in field theory, the procedure is much the same. We begin by the classical solution, expanding the energy functional about it and quantizing the leading order corrections. The simple harmonic analogous solutions are called mesons in quantum field theory, or quasiparticles in condensed matter. Of course, the mesons or quasiparticles also exist as excitations in the ground state, or vacuum state. Thus, quantization of the soliton is performed by accounting for the spectrum shift in the quasiparticle excitations and imposing commutation relations for the soliton position and momentum operators.

For a good introduction on the quantization of solitons from the quantum field theorist's point of view, see the book of Rajaraman⁵⁰ or the review articles of Coleman⁶ or Rajaraman⁴⁹.

Recall, however, that the soliton is a spontaneously broken symmetry solution: in has chosen an arbitrary point in space about which to center. The Goldstone theorem¹⁹ predicts a gapless boson mode restoring this broken symmetry. This causes divergences if we consider the next order semiclassical expansion of the quantized soliton, because of zero energy denominators that appear.

An analogous situation for a simple particle is when the potential is completely flat. To all orders we find zero frequencies when expanding the potential. This is because all points are degenerate and the particle must randomly choose among them. In the quantum version, we find that the particle is no longer an eigenvalue of position at all, but rather of momentum, in the form of a plane wave.

For the soliton, the Goldstone mode is dealt with in essentially the same way. For each broken symmetry, the quantized soliton has an associated momentum which is a good quantum number. For example, if the soliton exists in a translationally invariant system, we would find it has a well defined momentum in the quantized version. This, incidentally, provides a systematic method for calculating the mass of the soliton.

The general methods for separating the Goldstone mode involve introducing a collective coordinate for each broken symmetry^{18,20,68}. Since the original system doesn't depend on these coordinates, the final expanded energy functional can only depend on their conjugate momenta.

The magnetic system of this thesis has a two dimensional translational symmetry broken by the introduction of a vortex. Thus, we promote the vortex center coordinates to collective coordinates to we obtain an effective action depending only on the associated conjugate momentum via a particle-like $\frac{p^2}{2m}$ term.

1.3.2 In condensed matter theory

In condensed matter, we are more specifically interested in the physical consequences of the quantized solitons, as opposed to their mere existence and basic properties. Shortly after the quantum field theorists developed the soliton quantization methods, Krumhansl and Schrieffer^{7,34} showed that one dimensional quantized solitons could be treated exactly as elementary excitations, in addition to the ever-present quasiparticles. To explain, suppose we've quantized a soliton in a translationally invariant system (of length L with minimum length scale l). In the most general case, we would find, in addition to the regular Goldstone mode, a finite number of quasiparticle modes localized to the soliton, interpretable as soliton excited states, followed by the usual continuum of extended quasiparticle excitations. Krumhansl and Schrieffer show that the total internal energy of the system can be simplified to

$$U = \left(\frac{L}{l} - N_b N_k^{tot} \right) k_B T + N_k^{tot} \left(E_k^0 + \frac{1}{2} k_B T + (N_b - 1) k_B T \right) \quad (1.1)$$

where N_b is the total number of localized quasiparticle states, including the translation symmetry-restoring Goldstone mode.

This represents the internal energy of a system with $\left(\frac{L}{l} - N_b N_k^{tot} \right)$ quasiparticle modes and N_k^{tot} particles of rest energy E_k^0 each having $\frac{1}{2} k_B T$ translational energy and thermal energy $k_B T$ for each of the $N_b - 1$ internal modes. The average number of particles N_k^{tot} forming the soliton is calculated using thermodynamic relations once we define a soliton chemical potential. See Currie et al.⁷ for more details of the complete thermodynamic description of the soliton as an ideal gas.

Quantum vortices were first considered by condensed matter theorists as early as the 1940's by Onsager⁴⁵ in superfluid helium. Feynman¹⁴ developed further the idea of these vortex lines to explain the dissipation mechanisms for a rotating superfluid and conjectured that they may also be responsible for the superfluid to normal fluid phase transition. Unfortunately, in 3D the problem is essentially unsolved, so that no details of a vortex driven phase transition have yet been developed.

In 2D, the problem is more tractable, and in the 1970's, Kosterlitz and Thou-




Figure 1.7: The two degenerate dimer states of $trans-(CH)_N$, polyacetylene.

less³³ detailed a phase transition due to the proliferation of dislocations. The theory applies equally to vortices. Below the transition, the free energy is minimized by maintaining the vortex-antivortex pairs bound; however, raising the temperature to the transition, the gain in entropy by unbinding the pairs balances the increase in energy.

In two dimensions, the energy of a dislocation or vortex diverges logarithmically in the system surface area,

$$E = E_0 \ln \frac{A}{A_0} \quad (1.2)$$

where $A_0 \sim a^2$ is the smallest area in the discrete system, where a denotes the lattice spacing.

The entropy associated with the dislocation also depends logarithmically on the area since there are approximately A/A_0 possible positions for it to center on,

$$S = k_B \ln \frac{A}{A_0} \quad (1.3)$$

where k_B is the Boltzmann constant. Since the energy and entropy depend on the size of the system in the same way, the free energy, $F = E - TS$, is dominated by the energy term at low temperatures so that the probability of an isolated dislocation in a large system is vanishingly small. At high temperatures, dislocations appear spontaneously as the entropy term takes over. The phase transition temperature can be roughly estimated as $T_c = E_0/k_B$.

In the late 1970's, very important new phenomena were discovered independently by the particle and condensed matter physicists. Jackiw and Rebbi³⁰ in considering the Dirac equation in the presence of a soliton found it had fermionic $\frac{1}{2}$ states; while, Su, Schrieffer and Heeger⁶⁴ were studying kinks in a coupled electron-phonon model for the quasi-1D conducting polyacetylene and found a neutral spin $\frac{1}{2}$ soliton state.

Restricting ourselves to the polyacetylene system, consider a one dimensional system of electrons in a tight-binding model interacting linearly with the lattice coordinate displacements (essentially, coupling the electrons and phonons). The




Figure 1.8: The band structure of polyacetylene, gapped due to the electron-phonon interactions. Note the two isolated electron states in the gaps are only in the presence of a kink.

Hamiltonian of this system is then

$$\begin{aligned}
 H = & \sum_{n=1}^N \left(\frac{p_n^2}{2m} + \frac{K}{2} (u_{n+1} - u_n)^2 \right) - t_0 \sum_{n=1, s=\pm\frac{1}{2}}^N \left(c_{n+1,s}^\dagger c_{n,s} + c_{n,s}^\dagger c_{n+1,s} \right) \\
 & + \alpha \sum_{n=1, s=\pm\frac{1}{2}}^N (u_{n+1} - u_n) \left(c_{n+1,s}^\dagger c_{n,s} + c_{n,s}^\dagger c_{n+1,s} \right) \quad (1.4)
 \end{aligned}$$

where u_n and p_n are the lattice coordinate displacements and their conjugate momenta, characterized by mass m and stiffness constant K . The electrons are denoted by creation/annihilation operators $c_{i,s}^\dagger$ and $c_{i,s}$ at site i with spin s , with hopping constant t_0 and coupling constant α with the lattice displacements.

The ground state of this system is doubly degenerate and spontaneously breaks reflection symmetry (this was predicted by Peierls⁴⁷ using mean-field approximation for any non-zero electron-phonon coupling). Figure 1.7 shows the two degenerate dimer states. As a consequence of the two-fold degeneracy, there exist the kink and antikink topological solitons connecting the degenerate ground states (see Figure 1.9—in actuality, the kink is spread over $\sim 14a$).

Su et al.⁶⁴ found that the kink had two states: a charged state, $Q = \pm e$, with spin $s = 0$, and a neutral state with spin $s = \pm\frac{1}{2}$. In addition, when the kink is in its neutral state, there is an $s = 0$ electron state in the middle of the gap (see Figure 1.8, note there are two states, one localized to the kink, the other to the antikink) formed by pulling $\frac{1}{2}$ a state per spin out of the Fermi sea.

The polyacetylene study introduced to condensed matter physics what the particle theorists independently introduced within a relativistic field theory: the existence of states with fractional charge. Although the $\frac{1}{2}$ charge is obscured by the doubling of degrees of freedom due to spin, the zero energy state is




Figure 1.9: A kink solution connecting the two degenerate dimer ground states, shown, left, on the linear polyacetylene chain, and, right, on the idealized chain with periodic boundary conditions.

still formed by drawing half an electronic state (of each spin). Furthermore, the spin-charge relations are also unusual: charged solitons are spinless while neutral solitons carry spin $\frac{1}{2}$.

1.3.3 Superfluid He^4

Returning our discussion specifically to vortices in condensed matter, quantum vortices were first proposed by Onsager⁴⁵ and developed more completely by Feynman¹⁴. A quantum vortex can be imagined as a regular fluid vortex with a cylindrical core shrunk down to atomic dimensions. The circulation of the vortex is quantized in units of h/m , where h is the Planck constant and m is the bare ^4He mass.

Describing the motion of superfluid vortices by making analogy to the motion of their parent fluid vortices was extremely successful. Early experiments by Hall and Vinen^{23,24} found that if they applied an impulsive force setting a superfluid vortex into motion the vortex underwent helical motion (resembling that of an electron drifting in a magnetic field). In general, such a force arises always when a body with a flow circulation around it moves through a liquid or gas as in, for example, Figure 1.10.

First noted in 1852 by Magnus when studying inaccuracies in the firing of cannon balls, the force responsible, named the Magnus force after its discoverer, can be explained in terms of the Bernoulli equation. The speed of the fluid is effectively lower on one side of the rotating body than the other (perpendicularly to the flow of the fluid, of course) so that the side with higher speed has lower pressure—thus the body experiences a force in that direction (see Figure 1.10). The Magnus force in a superfluid is written

$$\mathbf{F}_M = \rho_s \kappa \times (\mathbf{v} - \mathbf{v}_s) \quad (1.5)$$

where ρ_s is the superfluid density, $\mathbf{v}$ is the vortex velocity and $\mathbf{v}_s$ is the asymp-




Figure 1.10: The equi-pressure lines of a fluid surrounding a rotating cylinder. The pressure differential top and bottom creates an upward force. The fluid flow is to the left.

otic superfluid velocity (affected, of course, by the vortex presence).

Hall and Vinen found the motion of their experimentally observed vortices could be explained with such a perpendicular Magnus force and an inertial mass of the order $\rho\xi^2$, where ρ is the fluid density and ξ is the vortex radius.

In addition, damping forces acting on the vortex were introduced with phenomenological parameters. The most general damping can act both longitudinal (as we are most accustomed to) and transverse to the vortex motion, expressible as

$$\mathbf{F}_d = D(\mathbf{v}_n - \mathbf{v}) + D'\hat{\kappa} \times (\mathbf{v}_n - \mathbf{v}) \quad (1.6)$$

where $\mathbf{v}_n$ denotes the normal fluid velocity, whose exact definition might vary from one formalism to another. Note that the transverse damping term has the same behaviour of the Magnus force (with potentially an additional force $\propto \mathbf{v}_n - \mathbf{v}_s$).

Although this heuristic description is very successful in explaining observed phenomena, the microscopic derivation of the various parameters is far less successful. There is considerable disagreement, especially in calculations of the transverse dissipation parameter.

An early calculation by Iordanskii^{26,27} revealed a transverse damping force, later termed the Iordanskii force, proportional to the normal fluid density

$$\mathbf{F}_I = \rho_n \kappa \times (\mathbf{v} - \mathbf{v}_n) \quad (1.7)$$

due to the scattering of phonons on the vortex. This entails an effective Magnus force with the superfluid density replaced by the total fluid density, $\rho_s \rightarrow \rho$, plus additional forces proportional to $\mathbf{v}_n - \mathbf{v}_s$.

In the early 1990's, Thouless, Ao and Niu^{2,67} (TAN) claimed that the transverse force was exactly the bare Magnus force of equation (1.5), at all temperatures

while accounting for the scattering of phonons. The force on the vortex line due to phonons is simply the variation of the phonon energy expectation with vortex position

$$\mathbf{F} = - \sum_{\alpha} f_{\alpha} \langle \psi_{\alpha} | \nabla_0 H | \psi_{\alpha} \rangle \quad (1.8)$$

where f_{α} denotes the occupation probability of the phonon state α . By expanding the phonon wavefunction to first order in vortex velocity using time-dependent perturbation theory, TAN were able to rewrite the force as an integral over the Berry phase associated with a closed loop around the vortex. Assuming no circulation in the normal fluid density, this reduces exactly to the zero temperature Magnus force.

The transverse force on the vortex line can also be expressed as the commutator of the x and y components of the total momentum operator

$$[P_x, P_y] = \int \int dx dy \left(\frac{\partial \psi^{\dagger}}{\partial x} \frac{\partial \psi}{\partial y} - \frac{\partial \psi^{\dagger}}{\partial y} \frac{\partial \psi}{\partial x} \right) \quad (1.9)$$

Applying Stokes' theorem, the integral over the cross-sectional area can be expressed instead as a line integral about the boundary of the one particle density matrix. TAN argue that this boundary may be extended very far from the vortex core so that contributions from localized phonon states at the vortex core do not influence the transverse force⁶⁷.

In opposition to Thouless, Sonin⁶¹ explained the transverse damping force via an analogous mechanism to the Aharonov-Bohm¹ effect of an electron passing a double slit in the presence of a magnetic vector potential (though in regions of no magnetic field). The electrons passing in one slit relative to the other experience a phase shift due to the vector potential term, causing a horizontal shift in the observed interference pattern. However, this entails a momentum transfer from the magnetic field source, here a conducting coil, to the electrons, transverse to the double slit screen, and thus a transverse force acting on the coil.

Similarly, quasiparticles passing above or below a moving vortex experience a relative Berry's phase shift⁴. A momentum transfer must occur between the vortex and quasiparticles, again, entailing a transverse damping force.

Sonin calculated the effective transverse force exactly in the form

$$\mathbf{F}_t = (\rho_s + \rho_n) \kappa \times (\mathbf{v} - \mathbf{v}_n) \quad (1.10)$$

so that the effective Magnus force is the regular Berry's phase result plus the Iordanskii force. The normal fluid velocity here is in the vicinity of the vortex and may differ from the asymptotic velocity due to viscous dragging of the normal fluid by the vortex motion²³.

One apparent source of disagreement, first noted by Sonin, is that the vortex undergoes oscillatory motion due to the passage of phonon quasiparticles. The

scattering calculations of Fetter¹³ and Demircan et al.⁹, which supported the TAN Berry's phase calculation, effectively held the vortex fixed by an external pinning potential, thereby nullifying the transverse damping force.

The transverse dissipation is not the only source of controversy. The effective mass itself of the quantized vortex has not been agreed upon. Initial estimates are based on the inertial mass of the circulating fluid, essentially, ρr_0^2 , with r_0 the radius of the vortex. In the quantum limit, the vortex radius shrinks down to atomic dimensions, or zero, so that the vortex mass tends to zero also.

Alternatively, as suggested by Duan and Leggett¹¹, the mass of the vortex must be proportional to

$$M_v \propto \frac{E_v}{v_0^2} \quad (1.11)$$

where M_v is the vortex mass, E_v is the stationary vortex energy, and v_0 is the velocity scale of the superfluid quasiparticles. This can be explained by purely dimensional arguments.

For a quasi-2D vortex, however, the stationary vortex energy is log divergent in the system cross-sectional area, as in (1.2), suggesting the effective mass is also log divergent, much larger than the vanishing estimate made earlier.

Clearly, the microscopic derivations of superfluid vortex dynamics has yet to firmly agreed upon. The variety of conflicting results suggests we re-examine the different methods used. Doing so in the simpler magnetic system is an aim of this thesis, though, unfortunately, a comprehensive study of the various methods could not entirely be undertaken. Rather, we calculate results here using regular perturbation theory, expanding in vortex velocity, and using Feynman-Vernon influence functionals¹⁷.

1.3.4 Magnetic vortices

Magnetic systems have received much attention for their variety of applications and their lucrative potential⁵², for example, in the market of magnetic memory. Vortices in magnetic systems are very easily observed and manipulated, for example using Brillouin light scattering⁴⁴ or magnetic force microscopy (MFM)⁵⁸.

Despite the ease of experimentally observing magnetic vortices, there have been relatively few microscopic derivations of the dynamics of vortices in magnetic systems. In fact, these derivations should be greatly simplified in a magnetic system; however, the resulting dynamics still possess many of the same strange aspects discussed with respect to superfluid vortices.

A magnetic vortex experiences a force transverse to its velocity, the *gyrotropic* force. This force acts exactly in the same manner as the Magnus force, however, has a different microscopic origin. It arises from a self induced Lorentz force, with the vortex vorticity acting as an analogous charge, while the out

of plane spins create an effective perpendicular magnetic field (this analogy is more fully developed in Chapter 3). Notably, this force is dependent on both topological indices (and is absent entirely for in-plane vortices for which the polarization is zero), as compared to the Magnus force dependent solely on the vortex circulation in a superfluid.

There are interactions with quasiparticles that may alter the effective gyrotropic force. However, there have been no attempts to describe a transverse damping force in a magnetic system. In fact, all descriptions of dissipation in a vortex system have focussed on calculating an average energy dissipation rate or have been phenomenological (except for the work of Slonczewski⁵⁹ which we describe in a moment).

The earliest theoretical work on two dimensional magnetic systems with vortices are adaptations of the work of Thiele^{65,66}. Thiele first introduced the gyrotropic force and dissipation dyadic acting on a magnetic domain wall in a three dimensional system. His dissipative force, however, was phenomenological employing a Gilbert damping parameter (the phenomenological damping parameter normally introduced into the so-called Landau-Lifshitz equations governing the magnetization dynamics).

In the early 1980's, applying the work of Thiele, Huber²⁵ and Nikiforov and Sonin⁴³ independently described the basic motion of a magnetic vortex. They calculated the gyrotropic force and phenomenological damping forces acting on a single vortex.

Slonczewski⁵⁹ shortly thereafter considered perturbations about a moving vortex, deducing an effective mass tensor. A collection of vortices behave strongly coupled and the inertial energy is not diagonal but rather must be expressed as $\frac{1}{2}M_{ij}v_i v_j$ where there is an implied double sum over the vortex indices i and j . He calculated the vortex dissipation via a frequency dependent imaginary mass term by studying the asymptotic behaviour of the lowest order vortex-magnon coupling. We will compare our dissipation results with those of Slonczewski in Chapter 4.

Scattering phase shifts have been calculated for a variety of planar magnetic systems^{21,48,51}. They were primarily interested in the thermodynamic behaviour³⁴ of such systems and searching for a vortex signature that could be measurable to verify a Kosterlitz-Thouless³³ transition. In fact, based on the modified spin correlations due to the presence of vortices, a central peak found in neutron-scattering experiments could be reproduced⁴⁰.

In a series of papers^{39,41,69}, Mertens et. al. modeled numerically the motion of a vortex pair assuming various boundary conditions. The ensuing motion was best reproduced assuming a non-Newtonian equation of motion which included a third time derivative of the vortex position.

We find just such a small third time derivative term in our influence functional

analysis. We compare our results with Mertens et. al. in section 4.3.4.

However, this is a misapplication of the collective coordinate formalism: each collective coordinate is meant to replace a continuous symmetry broken by the vortex. In a planar system, a vortex breaks the two dimensional translational symmetry allowing the introduction of a two dimensional center coordinate only.

There has been no work yet to find effective damping forces acting dynamically on a magnetic vortex. In this thesis, we calculate these forces assuming an averaged motion of the perturbing magnons.

1.4 Easy plane insulating ferromagnet

We study an insulating plane of spins, that is, fixed on their lattice sites, ferromagnetically coupled, lying preferentially in the plane. The order parameter of the easy plane ferromagnet lies on the unit sphere but with an energy barrier at both the north and south poles. There are hence topological solitons spontaneously breaking the ground state symmetry, the continuous in-plane symmetry, and, at some energy cost to restore continuity, twisting out of plane to break the discrete up/down symmetry. There are also discontinuous vortices lying entirely in the plane as found in the XY model.

We noted in the symmetry breaking discussion that a vortex lying in this order parameter space does not have topological stability. This however is for a completely degenerate sphere. Here, there is an energy barrier for paths to cross the two poles so that any homotopy of a vortex to a point would require passing a macroscopic number of spins through this energy barrier. The vortex thus has approximate topological stability, unless the anisotropy becomes vanishingly small.

The energy of a general state $\{\mathbf{S}_i\}$ of this lattice is

$$E = -\frac{1}{2} \sum_{i,j} J_{ij} \mathbf{S}_i \cdot \mathbf{S}_j + \sum_i K S_{iz}^2 \quad (1.12)$$

where the indices extend over all lattice points in the 2D lattice. The first term is the exchange term and is approximated by including nearest neighbour interactions only, negative to ensure ferromagnetic coupling,

$$-\frac{1}{2} \sum_{i,j} J_{ij} \mathbf{S}_i \cdot \mathbf{S}_j \rightarrow -\frac{1}{2} \sum_{\langle i,j \rangle} J \mathbf{S}_i \cdot \mathbf{S}_j$$

where $\langle i,j \rangle$ denotes nearest neighbour pairs. For simplicity, we've assumed a constant exchange parameter J . The second term enforces the easy plane anisotropy, where K is the anisotropy parameter (for $S > 1/2$).

Since we are interested in the low energy behaviour, we eliminate the short length scale fluctuations by describing the system in a continuum approximation. Instead of a spin $\mathbf{S}_i$ at site i , we now have a spin field $\mathbf{S}(\mathbf{r})$. Sums are replaced by integrals over space. For instance, the anisotropy term becomes

$$\sum_i K S_{iz}^2 \rightarrow \int d^2r \tilde{K} S_z^2(\mathbf{r})$$

and the exchange term becomes

$$-\frac{1}{2} \sum_{\langle i,j \rangle} J \mathbf{S}_i \cdot \mathbf{S}_j \sim \frac{1}{4} \sum_{\langle i,j \rangle} J (\mathbf{S}_i - \mathbf{S}_j) \cdot (\mathbf{S}_i - \mathbf{S}_j) \rightarrow \frac{1}{2} \int d^2r \tilde{J} (\nabla \mathbf{S})^2$$

where adding the constant S^2 terms doesn't affect the dynamics. Note that $(\nabla \mathbf{S})^2 = (\nabla S_x)^2 + (\nabla S_y)^2 + (\nabla S_z)^2$. The redefined constants are given by $\tilde{J} = J/2$ and $\tilde{K} = K/a^2$, noting that we use new dimensions for an anisotropy density. From here on, we drop the tildes and simply use J and K for the continuum versions of the exchange and anisotropy parameters.

The Hamiltonian describing the system is then written

$$H = S^2 \int d^2r \left(\frac{J}{2} (\nabla \theta)^2 + \sin^2 \theta \left(\frac{J}{2} (\nabla \phi)^2 - K \right) \right) \quad (1.13)$$

where the spin field is expressed in angular coordinates, $\mathbf{S} = S (\sin \theta \cos \phi, \sin \theta \sin \phi, \cos \theta)$.

As explained in Appendix A, ϕ and $-S \cos \theta$ are conjugate variables in the discrete lattice so that the Lagrangian can be expressed in the continuum limit, $\sum \rightarrow \int d^2r/a^2$ where a is some lattice spacing length scale,

$$\mathcal{L} = S \int \frac{d^2r}{a^2} \left(-\cos \theta \dot{\phi} - \frac{c}{2} \left((\nabla \theta)^2 + \sin^2 \theta \left((\nabla \phi)^2 - \frac{1}{r_v^2} \right) \right) \right) \quad (1.14)$$

where we've defined the speed scale c/r_v with $c = SJa^2$ and the length scale $r_v = \sqrt{J/2K}$.

Using Hamilton's equations (A.5) or the Euler-Lagrange equation (A.2), we find the equations of motion

$$\begin{aligned} \frac{1}{c} \frac{\partial \phi}{\partial t} &= -\frac{\nabla^2 \theta}{\sin \theta} + \cos \theta (\nabla \phi)^2 - \frac{1}{r_v^2} \cos \theta \\ \frac{1}{c} \frac{\partial \theta}{\partial t} &= \sin \theta \nabla^2 \theta + 2 \cos \theta \nabla \theta \cdot \nabla \phi \end{aligned} \quad (1.15)$$

There are two families of elementary excitations: the perturbative spin waves, or magnons, and the vortices. The vortices have two forms: the so-called in-plane

solutions with polarization 0, and the out-of-plane solutions with polarization ± 1 . The treatment in this thesis considers explicitly the out-of-plane solutions, however, setting the polarization to 0 recovers the results for the in-plane solutions. The out of plane spin behaviour cannot be solved analytically; however, the core and far field asymptotic limits suffice for obtaining general results.

The spin waves are small amplitude oscillations about the ferromagnetic ground state or about a vortex state, in both cases with an ungapped spectrum. The difference in the two spectra can be attributed to the vortex presence and yields an effective zero point energy to the quantized vortex. The equations of motion for the vacuum magnons are modified to the equations of motion of magnons in the presence of a vortex. The additional terms are interpreted as the magnon-vortex interaction terms.

There is a one magnon coupling with the vortex velocity. Normally, considering a central system coupled to perturbative ‘bath’ modes, we find to lowest order a one magnon coupling with the vortex field. There is no such coupling here because the vortex is itself a minimum action solution of the same system in which the magnons arise. Thus, there are no first order variational terms. This assumes, however, that the vortex profile is unchanging in time. Allowing it to move about the system introduces a first order coupling between the vortex velocity and the magnons.

There is also a two magnon coupling affecting the magnon energy with long range effects. This term scatters the magnon modes and hence alters their zero point energy. We attribute this shift instead to the quantized vortex state. This two magnon coupling has other dissipative effects and energy shifts that are not treated in this thesis.

We first review the basic characteristics of the vacuum magnon modes and the vortex solutions. The gyrotropic and inter-vortex forces are found immediately by expanding the Lagrangian about a many vortex solution.

We then examine the effects of the various couplings between magnons and vortices. The one magnon coupling can be interpreted as small vortex deformations when moving at velocity $\mathbf{V}$ or, alternatively, as a single magnon scattering event. The second order perturbation energy correction of this one magnon coupling goes as V^2 and is thus interpretable as an inertial energy, from which we can deduce an effective vortex mass. There is an additional imaginary energy shift, or a dissipation, from this coupling.

The two magnon scattering term has a zero point energy shift and other magnon occupation dependent energy shifts. We do not retain higher order scattering terms, keeping only one magnon couplings, although they may indeed contribute more significantly to the vortex dissipation^{12,62}.

The dynamical effect of the one magnon coupling is examined fully in the Feynman-Vernon influence functional formalism¹⁷. The two sub-systems are as-

sumed initially non-interacting with the magnons in thermal equilibrium. They are thereafter allowed to interact, the magnons generally shifting out of equilibrium. The effect of the magnons is then averaged over by tracing out their degrees of freedom. This yields, in an averaged way, the effect of the magnons on the vortex motion. As found in perturbation theory, the one magnon coupling is responsible for two new terms in the vortex effective action: an inertial energy term and a damping force term.

In addition to the usual longitudinal damping force, we find a transverse damping force reminiscent of the Iordanskii force in superfluid helium. Such a term has not before been suggested in a magnetic system. The damping forces possess memory effects—that is, they depend on the previous motion of the vortices.

For a collection of vortices, we find that their particle-like properties are not independent. They have mixed inertial terms such as $\frac{1}{2}M_{ij}v_i v_j$ and damping forces due to the motion of one vortex acting on another.

Next, we review the basics of the two elementary excitations, first the magnons and after the vortices.

Chapter 2

Magnons

The plane of spins with easy plane anisotropy has a degenerate ground state. The spins are ferromagnetically coupled and thus prefer to align, however, they may choose to align along any direction in the plane—an example of spontaneously broken symmetry. The Goldstone theorem predicts that there should then exist boson quasiparticle excitations that are not gapped. In this system, these Goldstone modes are the small amplitude, or perturbative, spin waves. When quantized, the excitations are termed magnons.

The magnon spectrum in the easy plane ferromagnet is ungapped, however due to the hard axis, the spectrum is not simply the regular ferromagnet spin wave spectrum $\omega(k) \propto k^2$. Instead we find a spectrum with reduced density of states near $\omega = 0$.

We begin by examining the small amplitude equations of motion satisfied by the magnons; thereby deriving the magnon spectrum and density of states. We calculate a few old results using spin path integrals as illustrative examples that we will need in later calculations. We derive the quantum propagator, a calculation following closely that of a simple harmonic oscillator. The quantum propagator is then manipulated to again reveal the magnon spectrum and, under a simple substitution, to yield the thermal equilibrium density matrix.

2.1 Magnon equations of motion

The magnons are the quasiparticle excitations of our system. As such, to describe their motion and properties, we expand in small deviations about the ferromagnetic in-plane ground state

$$\begin{aligned}\vartheta &= \theta - \pi/2 \\ \varphi &= \phi\end{aligned}\tag{2.1}$$

where we've chosen the ground state $\phi = 0$ amongst the continuum of ground states without loss of generality. The complete system Lagrangian in terms of these perturbing variables φ and ϑ becomes

$$\mathcal{L}_m = S \int \frac{d^2r}{a^2} \left(\dot{\varphi} \vartheta - \frac{c}{2} \left(-\varphi \nabla^2 \varphi - \vartheta \nabla^2 \vartheta + \frac{\vartheta^2}{r_v^2} \right) \right) \quad (2.2)$$

where J is the exchange constant and K is the anisotropy constant, a is the lattice spacing, $c = SJa^2$ and $r_v^2 = \frac{J}{2K}$. The conjugate momentum is now $S\vartheta$, the linearized version of $-S \cos \theta$. We essentially expand the Lagrangian to second order perturbations to obtain a simple harmonic-like Lagrangian. Consequently, many calculations to come here mimic very closely those for a simple harmonic oscillator.

Varying (2.2) with respect to φ and ϑ yields the magnon equations of motion

$$\begin{aligned} \frac{1}{c} \frac{\partial \varphi}{\partial t} &= -\nabla^2 \vartheta + \frac{1}{r_v^2} \vartheta \\ \frac{1}{c} \frac{\partial \vartheta}{\partial t} &= \nabla^2 \varphi \end{aligned} \quad (2.3)$$

Alternatively, we could have linearized the system equations of motion, (1.15), directly with identical results.

The analysis proceeds in a plane-wave expansion. This system of equations can be solved by Fourier transforming so that $\nabla^2 \varphi \rightarrow -k^2 \varphi_k$ and $\nabla^2 \vartheta \rightarrow -k^2 \vartheta_k$. Assuming harmonic time dependence, the eigenvalues of the equations of motion yield the magnon spectrum

$$\omega(k) = ckQ \quad (2.4)$$

where $Q = \sqrt{k^2 + \frac{1}{r_v^2}}$. The spectrum is not gapped (note the overall factor of k), a reflection of the continuous degeneracy of ground states. However, the density of states goes as $\frac{Q}{k^2 + Q^2}$ remaining finite as $\omega \rightarrow 0$ in comparison to the isotropic ferromagnet with density of states $\frac{1}{k}$ diverging for zero frequency. The two systems are compared in Figure 2.1.

Alternatively, Fourier transforming the magnon Hamiltonian directly via

$$\begin{aligned} \varphi(\mathbf{r}) &= \frac{a^2}{(2\pi)^2} \int d^2k e^{-i\mathbf{k} \cdot \mathbf{r}} \varphi_{\mathbf{k}} \\ \vartheta(\mathbf{r}) &= \frac{a^2}{(2\pi)^2} \int d^2k e^{-i\mathbf{k} \cdot \mathbf{r}} \vartheta_{\mathbf{k}} \end{aligned} \quad (2.5)$$

diagonalizes $H_m = \int \frac{d^2r}{a^2} S \dot{\varphi} \dot{\vartheta} - \mathcal{L}_m$ to

$$H_m = \frac{Sc}{2} \int \frac{d^2ka^2}{(2\pi)^2} (k^2 \varphi_{\mathbf{k}} \varphi_{-\mathbf{k}} + Q^2 \vartheta_{\mathbf{k}} \vartheta_{-\mathbf{k}}) \quad (2.6)$$




Figure 2.1: A comparison of the easy plane magnon spectrum and density of states with the regular isotropic ferromagnet.

To quantize the magnons, we impose the commutation relations between the conjugate variables $\varphi_{\mathbf{k}}$ and $S\vartheta_{\mathbf{k}}$

$$[S\vartheta_{\mathbf{k}}, \varphi_{\mathbf{k}'}] = -i\hbar(2\pi)^2 \frac{\delta^2(\vec{k} - \vec{k}')}{a^2} \quad (2.7)$$

We diagonalize the system now via the transformation to creation/annihilation operators

$$\begin{aligned} a_{\mathbf{k}} &= \sqrt{\frac{Sk}{2\hbar Q}} \left(\varphi_{\mathbf{k}} + \frac{iQ}{k} \vartheta_{-\mathbf{k}} \right) \\ a_{\mathbf{k}}^{\dagger} &= \sqrt{\frac{Sk}{2\hbar Q}} \left(\varphi_{-\mathbf{k}} - \frac{iQ}{k} \vartheta_{\mathbf{k}} \right) \end{aligned} \quad (2.8)$$

normalized such that $[a_{\mathbf{k}}, a_{\mathbf{k}'}^{\dagger}] = (2\pi)^2 \frac{\delta^2(\mathbf{k}-\mathbf{k}')}{a^2}$. Substituting for $\varphi_{\mathbf{k}}$ and $\vartheta_{\mathbf{k}}$ in terms of $a_{\mathbf{k}}$ and $a_{\mathbf{k}}^{\dagger}$ into the Fourier transformed Hamiltonian gives after some manipulation

$$\begin{aligned} H &= \int \frac{a^2 d^2 k}{(2\pi)^2} \frac{\hbar\omega_k}{2} \left(a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} + a_{\mathbf{k}} a_{\mathbf{k}}^{\dagger} \right) \\ &= \int \frac{a^2 d^2 k}{(2\pi)^2} \hbar\omega_k \left(a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} + \frac{1}{2} \right) \end{aligned} \quad (2.9)$$

where ω_k is again the magnon dispersion relation (2.4).

We interpret the operators $a_{\mathbf{k}}^{\dagger}$ and $a_{\mathbf{k}}$ exactly as for the simple harmonic oscillator creation/annihilation operators. The combination $a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}}$ is thus the

magnon number operator $n_{\mathbf{k}}$ and the spectrum has energy $\hbar\omega_{\mathbf{k}}$ for each of the $n_{\mathbf{k}}$ magnons plus an additional *zero-point energy* $\frac{1}{2}\hbar\omega_{\mathbf{k}}$ for each wavevector $\mathbf{k}$.

Notice throughout that we associate factors of a^2 to the spacial and frequency integration measures to keep them dimensionless. This is consistent since the integrals replace sums appearing in the original discrete system.

2.2 Quantum propagator

The quantum propagator is an operator describing the time evolution of a quantum state. Although the vacuum propagator of the magnons is not needed for future calculations in this thesis, its calculation offers a simple application of spin path integration in our easy-plane ferromagnet. With only slight modifications to this derivation, that is with the addition of a perturbing term, or forcing term, we obtain the quantum propagator for magnons in the presence of a vortex. We must save this calculation for later after we've derived the appropriate forcing term.

Suppose initially we know the state of the system of magnons which can be represented in the φ basis. To find the state of the system at a later time, T ,

$$\psi(\varphi, T) = \int d\varphi' K(\varphi, T; \varphi', 0) \psi(\varphi', 0) \quad (2.10)$$

where

$$K(\varphi, T; \varphi', 0) \equiv \langle \varphi | \exp - \frac{iHT}{\hbar} | \varphi' \rangle \quad (2.11)$$

is the quantum propagator expressible as a path integral (see Appendix C)

$$K(\varphi, T; \varphi', 0) = \int_{\varphi'}^{\varphi} \mathcal{D}[\varphi(\mathbf{r}, t), \vartheta(\mathbf{r}, t)] \exp \left(\frac{i}{\hbar} \int_0^T dt \mathcal{L}_m[\varphi, \vartheta] \right) \quad (2.12)$$

and where $\mathcal{S}_m = \int_0^T dt \mathcal{L}_m$ is the action with the Lagrangian $\mathcal{L}_m$ given in (2.2).

Before proceeding with the semiclassical approximation—here exact since we have no terms of higher order than quadratic—first Fourier transform to diagonalize the problem in k -space. Introducing the Fourier pairs of φ and ϑ , (2.5), the Lagrangian becomes

$$\mathcal{L}_m = S \int \frac{a^2 d^2 k}{(2\pi)^2} \left(\dot{\varphi}_{\mathbf{k}} \vartheta_{-\mathbf{k}} - \frac{c}{2} (k^2 \varphi_{\mathbf{k}} \varphi_{-\mathbf{k}} + Q^2 \vartheta_{\mathbf{k}} \vartheta_{-\mathbf{k}}) \right) \quad (2.13)$$

The path integration measure is now the product of these Fourier coefficients

$$\mathcal{D}[\varphi(\mathbf{r}, t), \vartheta(\mathbf{r}, t)] \rightarrow \prod_{\mathbf{k}} d\varphi_{\mathbf{k}}(t) d\vartheta_{\mathbf{k}}(t)$$

Now to find the classical contribution, the equations of motion arising from this Lagrangian are

$$\begin{pmatrix} ck^2 & \frac{\partial}{\partial t} \\ -\frac{\partial}{\partial t} & cQ^2 \end{pmatrix} \begin{pmatrix} \varphi_{\mathbf{k}}^{cl} \\ \vartheta_{\mathbf{k}}^{cl} \end{pmatrix} = 0 \quad (2.14)$$

The general solution with boundary conditions $\varphi_{\mathbf{k}}(0) = \varphi'_{\mathbf{k}}$ and $\varphi_{\mathbf{k}}(T) = \varphi_{\mathbf{k}}$ is

$$\begin{pmatrix} \varphi_{\mathbf{k}}^{cl} \\ \vartheta_{\mathbf{k}}^{cl} \end{pmatrix} = \frac{\varphi_{\mathbf{k}}}{\sin \omega_k T} \begin{pmatrix} \sin \omega_k (T-t) \\ \frac{k}{Q} \cos \omega_k t \end{pmatrix} + \frac{\varphi'_{\mathbf{k}}}{\sin \omega_k T} \begin{pmatrix} \sin \omega_k (T-t) \\ -\frac{k}{Q} \cos \omega_k (T-t) \end{pmatrix} \quad (2.15)$$

where $\omega_{\mathbf{k}} = ckQ$ as before.

Substituting the classical solution back into the action, after some simplification, yields the classical contribution to the action

$$\mathcal{S}_m^{cl} = \int \frac{a^2 d^2 k}{(2\pi)^2} \frac{Sk}{2Q \sin \omega_k T} ((\varphi'_{\mathbf{k}} \varphi'_{-\mathbf{k}} + \varphi_{\mathbf{k}} \varphi_{-\mathbf{k}}) \cos \omega_k T - 2\varphi_{\mathbf{k}} \varphi'_{-\mathbf{k}}) \quad (2.16)$$

To evaluate the effect of quantum fluctuations, we solve the relevant Jacobi equation (adapted for a spin path integral as described in Appendix C)

$$\begin{pmatrix} ck^2 & \frac{\partial}{\partial t} \\ -\frac{\partial}{\partial t} & cQ^2 \end{pmatrix} \begin{pmatrix} \varphi(t) \\ \vartheta(t) \end{pmatrix} = 0 \quad (2.17)$$

with initial conditions $\varphi(0) = 0$ and $S\vartheta(t) = 1$. The determinant of the fluctuations is then given by $ix(T) = iSQ/k \sin \omega_k T$ for each k . Combined with the prefactors in the path integration measure $S/\hbar$, we find that the Gaussian integral over fluctuations yields the prefactor

$$\sqrt{\frac{Sk}{2\pi i \hbar Q \sin \omega_k T}} \quad (2.18)$$

Assembling the various pieces, the propagator of the unperturbed magnons is

$$K(\varphi, T; \varphi', 0) = \prod_{\mathbf{k}} \sqrt{\frac{Sk}{2\pi i \hbar Q \sin \omega_k T}} \exp \left(\int \frac{a^2 d^2 k}{(2\pi)^2} \frac{iSk}{2\hbar Q \sin \omega_k T} ((\varphi'_{\mathbf{k}} \varphi'_{-\mathbf{k}} + \varphi_{\mathbf{k}} \varphi_{-\mathbf{k}}) \cos \omega_k T - 2\varphi_{\mathbf{k}} \varphi'_{-\mathbf{k}}) \right) \quad (2.19)$$

where $\varphi_{\mathbf{k}}$ and $\varphi'_{\mathbf{k}}$ are the Fourier components of the boundary functions $\varphi(\mathbf{r})$ and $\varphi'(\mathbf{r})$.

2.2.1 Spectrum via tracing over the propagator

By manipulating the propagator, we can recover the magnon spectrum. To explain, consider first the propagator of a single particle starting from position q_0 at time 0 and going to position q_T at time T

$$K(q_T, T; q_0, 0) = \langle q_T | \exp -\frac{iHT}{\hbar} | q_0 \rangle \quad (2.20)$$

Taking the trace of this operator, i.e. set $q_T = q_0$ and integrate over the end-point q_0 of the periodic orbit, we find

$$\begin{aligned} G(T) &= \int_{-\infty}^{\infty} dq_0 \langle q_0 | \exp -\frac{iHT}{\hbar} | q_0 \rangle \\ &= \int_{-\infty}^{\infty} dq_0 \sum_n \langle q_0 | \xi_n \rangle \exp -\frac{iE_n T}{\hbar} \langle \xi_n | q_0 \rangle \\ &= \sum_n \exp -\frac{iE_n T}{\hbar} \end{aligned} \quad (2.21)$$

where $\{\xi_n\}$ denote a complete orthonormal set of eigenstates of H . Using the normalization condition of these ξ_n then yields the excitation spectrum of the Hamiltonian.

The trace of the propagator (2.19) thus provides another means to find the excitation spectrum. Set $\varphi_{\mathbf{k}} = \varphi'_{\mathbf{k}}$ and integrate over each Fourier coefficient

$$G_m(T) = \prod_{\mathbf{k}} \int d\varphi_{\mathbf{k}} \sqrt{\frac{Sk}{2\pi i \hbar Q \sin \omega_k T}} \exp \left(\frac{iSk}{\hbar Q \sin \omega_k T} \varphi_{\mathbf{k}} \varphi_{-\mathbf{k}} (\cos \omega_k T - 1) \right)$$

For ease of notation, assume that the product function applies to everything to the right of it, notably implying an integration over $\mathbf{k}$ within the exponential.

Performing the Gaussian integrals

$$G_m(T) = \prod_{\mathbf{k}} \sqrt{\frac{Sk}{2\pi i \hbar Q \sin \omega_k T}} \sqrt{\frac{\pi \hbar Q \sin \omega_k T}{-iSk (\cos \omega_k T - 1)}}$$

and noting that $\cos \omega_k T - 1 = -2 \sin^2 \left(\frac{\omega_k T}{2} \right)$, this reduces to

$$\begin{aligned}
G_m(T) &= \prod_{\mathbf{k}} \frac{1}{2i \sin\left(\frac{\omega_{\mathbf{k}} T}{2}\right)} \\
&= \prod_{\mathbf{k}} e^{-i\omega_{\mathbf{k}} T/2} \frac{1}{1 - e^{-i\omega_{\mathbf{k}} T}} \\
&= \prod_{\mathbf{k}} \sum_{n=0}^{\infty} e^{-i(n+\frac{1}{2})\omega_{\mathbf{k}} T} \\
&= \sum_{\{n_{\mathbf{k}}\}} e^{-i \sum_{\mathbf{k}} (n+\frac{1}{2})\omega_{\mathbf{k}} T} \tag{2.22}
\end{aligned}$$

where $\{n_{\mathbf{k}}\}$ denotes a set of integers $n_{\mathbf{k}}$. Thus, comparing with equation (2.21), we find the excitation spectrum $\sum_{\mathbf{k}} \hbar\omega_{\mathbf{k}}(n + \frac{1}{2})$, as expected.

This method of recovering the excitation spectrum is of course only useful in the special case that G can be cast into this final form. Nonetheless, by taking the limit $T \rightarrow 0$, the ground state term dominates the summation so that we can always at least find the ground state energy.

2.3 Thermal equilibrium density matrix

2.3.1 Magnon density matrix

A quantum state is represented by a wavefunction $\psi(\mathbf{r}, t)$. Generally, this state is a superposition of the system energy eigenstates, $\{\xi_i\}$. For example

$$\psi(\mathbf{r}, t) = \sum_i c_i \xi_i \tag{2.23}$$

The probability of finding the i th eigenstate upon measurement is c_i^2 and, by conservation of probability, $\sum c_i^2 = 1$. This is a pure quantum state. Alternatively, a system may be a statistical mixture of eigenstates. In that case, the quantum state isn't expressible as in (2.23), but, rather, is described by a set of probabilities p_i of finding the system in eigenstate ξ_i upon measurement.

We may have a pure quantum state describing the entire interacting system, which to some extent is the entire universe. Of course, we may then only be interested in a small subsystem within the whole. We wish to describe its quantum state only in terms of the subsystem coordinates.

The density matrix is a notation for describing a quantum state, necessitated by statistical mixtures such as a thermal equilibrium state, or entangled states of two sub-systems for which each individual system must be described by a density matrix even though the complete system may be in a pure state. As the

name implies, we express the quantum state by a matrix describing the density of the subsystem or mixture in terms of its eigenstates or coordinates. More specifically, for a pure state, the density matrix is

$$\rho_{ij} = c_i c_j \quad (2.24)$$

where the c_i are the coefficients in (2.23). For a mixture,

$$\rho_{ij} = \delta_{ij} p_i \quad (2.25)$$

where p_i are again the probabilities of finding the system in state ξ_i .

Supposing we have a pure quantum state, we can write the density matrix in the coordinate basis

$$\begin{aligned} \rho(\mathbf{x}, \mathbf{x}') &= \sum_{ij} \xi_i(\mathbf{x}) \rho_{ij} \xi_j^*(\mathbf{x}') \\ &= \psi(\mathbf{x}) \psi^*(\mathbf{x}') \end{aligned}$$

The vector $\mathbf{x}$ is broken into the coordinates of interest $\tilde{\mathbf{x}}$ and remaining coordinates $\mathbf{q}$ such that $\mathbf{x} = (\tilde{\mathbf{x}}, \mathbf{q})$. The *reduced* density matrix for the subsystem of interest is found by tracing out the uninteresting degrees of freedom

$$\tilde{\rho}(\tilde{\mathbf{x}}, \tilde{\mathbf{x}}') = \int d\mathbf{q} \rho(\tilde{\mathbf{x}}, \mathbf{q}; \tilde{\mathbf{x}}', \mathbf{q})$$

This effectively averages the effects of the external system. For a pure state, $\text{tr} \rho^2 = 1$. It can be shown that $\text{tr} \rho^2$ is maximal when the ensemble is pure; for a mixed ensemble $\text{tr} \rho^2$ is a positive number less than one.

A quantum system in thermal equilibrium has its eigenstates populated with probabilities given by the Boltzmann weighting factor $e^{-\beta E_i}$. The thermal density matrix in the coordinate basis is

$$\rho(\mathbf{x}, \mathbf{x}') = \sum_i \xi_i(\mathbf{x}) \xi_i^*(\mathbf{x}') e^{-\beta E_i} \quad (2.26)$$

This should be normalized by the partition function $\sum e^{-\beta E_i}$; however, we will omit it for ease of notation.

But this form is extremely similar to the quantum propagator when also expressed in this basis

$$\begin{aligned} K(\mathbf{x}, T; \mathbf{x}', 0) &= \sum_i \xi_i(\mathbf{x}) \xi_i^*(\mathbf{x}') e^{-i/\hbar H E_i} \\ &= \langle \mathbf{x} | e^{-i/\hbar H E_i} | \mathbf{x}' \rangle \end{aligned}$$

Under the substitution $T \rightarrow -i\hbar\beta$, in fact, we recover the thermal density matrix, though unnormalized. See Appendix A.1 for formal details of this substitution.

Noting this imaginary time correspondence between the quantum propagator and the density matrix in thermal equilibrium, we make the substitution $T \rightarrow -i\hbar\beta$ in the quantum propagator

$$K(\varphi, T; \varphi', 0) = \prod_{\mathbf{k}} \sqrt{\frac{Sk}{2\pi i\hbar Q \sin \omega_k T}} \exp \left(\int \frac{a^2 d^2 k}{(2\pi)^2} \frac{iSk}{2Q \sin \omega_k T} \right. \\ \left. ((\varphi'_{\mathbf{k}} \varphi'_{-\mathbf{k}} + \varphi_{\mathbf{k}} \varphi_{-\mathbf{k}}) \cos \omega_k T - 2\varphi_{\mathbf{k}} \varphi'_{-\mathbf{k}}) \right)$$

to obtain the thermal equilibrium density matrix

$$\rho(\varphi, \varphi') = \prod_{\mathbf{k}} \sqrt{\frac{Sk}{2\pi \hbar Q \sinh \hbar \omega_k \beta}} \exp \left(- \int \frac{a^2 d^2 k}{(2\pi)^2} \frac{Sk}{2Q \sinh \hbar \omega_k \beta} \right. \\ \left. ((\varphi'_{\mathbf{k}} \varphi'_{-\mathbf{k}} + \varphi_{\mathbf{k}} \varphi_{-\mathbf{k}}) \cosh \hbar \omega_k \beta - 2\varphi_{\mathbf{k}} \varphi'_{-\mathbf{k}}) \right) \quad (2.27)$$

This corresponds to the magnons being excited such that a state with energy E_k is measured with probability weighting given by the Boltzmann factor, $e^{-i\beta E_k}$.

2.4 Summary

In summary, the easy-plane magnons perturbing the vacuum ground state have the spectrum

$$\omega(k) = ckQ$$

where $Q = \sqrt{k^2 + \frac{1}{r_v^2}}$, $c = SJa^2$ and $r_v^2 = \frac{J}{2K}$.

We calculated the real time propagator of these magnons and consequently, making use of the imaginary time path integral of the density matrix, also the thermal equilibrium density matrix. The propagator is extremely similar to that of a simple harmonic oscillator. In fact, under the substitution $\frac{Sk}{Q} \rightarrow m\omega$ the magnon propagator becomes identical to that of the simple harmonic oscillator.

Next, we examine the vortex excitations.

Chapter 3

Vortices

The easy plane ferromagnet admits two families of elementary excitations. In the last chapter, we reviewed the perturbative excitations, the magnons. Now, we review the other elementary excitations, the non-perturbative vortices.

Although the out-of-plane spin behaviour cannot be described analytically, we present the asymptotic behaviour which is sufficient for getting leading order results. By superposing many vortex solutions, we expand the action to reveal an inter-vortex Coulomb-like force. The analogy is complete with the correspondence of $4\pi\epsilon_0 q_i$ with electronic charge in Coulombs.

The dynamic term “ $p\dot{q}$ ” in the action is re-expressed describing a gyrotropic force (analogous to the Lorentz force) or, alternatively, as an effective dynamic term in terms of vortex coordinates, $\mathbf{P} \cdot \dot{\mathbf{X}}$, where the momentum term is a vector potential. This is analogous to a charge in a magnetic field for which the momentum is modified by the magnetic field vector potential. In this formalism, the corresponding vector potential describes an effective perpendicular magnetic field $\mathbf{B} = \frac{S^2 J}{4\epsilon_0 r_v} p_i \hat{z}$.

We briefly present different possible two-vortex motions: depending on the relative sign of $p_i q_i$, the pair execute parallel motion (for opposite signs) or co-orbital motion (for like signs). This basic motion is perturbed by introducing an inertial mass term. Finally, the zero point energy shift of the two magnon coupling is examined in a Born approximation. This approximation is found to be sufficient for the continuum of magnons; however, there exist translation modes localized to the vortex core. These will be reconsidered in the next chapter using collective coordinates.

The system has two symmetries: a continuous in-plane symmetry and a discrete up-down symmetry. The vortices are thus characterized by two topological indices, the vorticity $q = \pm 1, \pm 2, \dots$, sometimes also called the winding number, and the polarization $p = 0, \pm 1$.

The $p = 0$ vortices are often separately considered, termed the in-plane vortices, while the $p \neq 0$ solutions are called the out-of-plane vortices. This separation, however, is unnecessary: allowing $p = 0, \pm 1$ in the following treatment recovers the proper results for both types of solutions.

Being non-perturbative solutions, the vortices satisfy the full, non-linear, equations of motion of the easy plane ferromagnet. Derived from the system Lagrangian

$$\mathcal{L} = S \int \frac{d^2r}{a^2} \left(-\dot{\phi} \cos \theta - \frac{c}{2} \left((\nabla \theta)^2 + \sin^2 \theta \left((\nabla \phi)^2 - \frac{1}{r_v^2} \right) \right) \right) \quad (3.1)$$

the equations of motion are

$$\begin{aligned} \frac{1}{c} \frac{\partial \phi}{\partial t} &= -\frac{\nabla^2 \theta}{\sin \theta} + \cos \theta (\nabla \phi)^2 - \frac{1}{r_v^2} \cos \theta \\ \frac{1}{c} \frac{\partial \theta}{\partial t} &= \sin \theta \nabla^2 \theta + 2 \cos \theta \nabla \theta \cdot \nabla \phi \end{aligned} \quad (3.2)$$

where J is the exchange constant and K is the anisotropy constant, a is the lattice spacing, $c = SJa^2$ and $r_v^2 = \frac{J}{2K}$.

The in-plane vortex can be described analytically. The spin configuration of this solution has $\phi_v = q\xi + \delta$ and $\theta_v = 0$. The parameter q is called the *vorticity* of the vortex, and δ is a phase that has little importance on the vortex dynamics[†]. We can solve for its energy within our continuum approximation, requiring both an infrared and ultraviolet cutoff,

$$E = S^2 \int d^2r \frac{J}{2} (\nabla \phi_v)^2 = S^2 J \pi q^2 \ln \frac{R_s}{a} \quad (3.3)$$

where R_s is the radial size of the system and a is a lower cutoff, the lattice spacing, required since the system is actually discrete (making $r \rightarrow 0$ unphysical). Note that this energy is independent of where the vortex center is within the circular integration region.

The out-of-plane solution is also characterized by its polarization; that is, the direction (up/down) that the spins twist out-of-plane. The spin configuration has the same polar angle dependence, $\phi_v = q\xi + \delta$, while the out-of-plane spin angle cannot be solved for analytically. The asymptotic behaviour is

$$\cos \theta_v = \begin{cases} 1 - c_1 \left(\frac{r}{r_v} \right)^2, & r \rightarrow 0; \\ c_2 \sqrt{\frac{r_v}{r}} \exp\left(-\frac{r}{r_v}\right), & r \rightarrow \infty. \end{cases} \quad (3.4)$$

where c_1 and c_2 are free constants that can be set by imposing appropriate continuity conditions.

Figure 3.1 shows the spin configuration of two simple out-of-plane vortices. This solution has the same leading order energy as the in-plane solution

[†]Note that this broken continuous symmetry entails the existence of gapless boson modes: the magnons.




Figure 3.1: Vortex spin configuration: left, a vortex with $q = -1$; right, a vortex with $q = 1$.

$$E_v = S^2 J \pi q^2 \ln \frac{R_s}{a} \quad (3.5)$$

Core corrections to the energy are finite and hence negligible in comparison to this log divergent contribution. In fact, in most that follows, the core will be ignored since it usually offers a finite contribution next to a log divergent one. A notable exception is the gyrotropic force that depends on the core behaviour via the core polarization. This is a differentiating feature of magnetic vortex dynamics from that of classical fluid or superfluid vortices where the analogous Magnus force depends only on the vortex circulation, the fluid analogue to the magnetic vorticity.

The motion of the in-plane vortex undergoes many of the same corrections. In fact, with the substitution $p \rightarrow 0$ the treatment here reduces to that of an in-plane vortex. The gyrotropic force disappears, however, all other forces and correction are polarization independent.

3.1 Force between vortices

Consider two vortices of vorticity q_1 and q_2 and polarization p_1 and p_2 well separated so that the only distortion in their profiles can be assumed to lie in the region between the two where their profiles are entirely in the plane. The spins in this middle ground are aligned in the plane with angle ϕ_{12} determined by the sum of spin angles (see Figure 3.2) given by each vortex independently

$$\phi_{12} = q_1 \chi(X_1) + q_2 \chi(X_2) \quad (3.6)$$

The out-of-plane component of the spin can be neglected here since we've as-




Figure 3.2: Two vortex spin configurations. Left, two vortices with $q = 1$; right, vortices with $q = 1$ and $q = -1$; both with no relative phase shift.

sumed that the vortex cores are widely separated and each core gives only a small correction.

The energy of the two vortex system is

$$E_{12} = \frac{S^2 J}{2} \int d^2 r \left((\nabla \theta_{12})^2 + \sin^2 \theta_v (\nabla \phi_{12})^2 + \frac{\cos^2 \theta_v}{r_v^2} \right) \quad (3.7)$$

which, except for regions within radius r_v of each vortex core, is dominated by the $(\nabla \phi_{12})^2$ term. Thus, neglecting core terms, the energy becomes

$$\begin{aligned} E_{12} &= \frac{S^2 J}{2} \int d^2 r (\nabla \phi_{12})^2 \\ &= \frac{S^2 J}{2} \int d^2 r \left(\frac{q_1 \hat{\phi}_1}{X_1} + \frac{q_2 \hat{\phi}_2}{X_2} \right)^2 \end{aligned} \quad (3.8)$$

As an illuminating trick to evaluating this integral, note that

$$\left(\frac{q_1 \hat{\phi}_1}{X_1} + \frac{q_2 \hat{\phi}_2}{X_2} \right)^2 = \left(\frac{q_1 \hat{\mathbf{X}}_1}{X_1} + \frac{q_2 \hat{\mathbf{X}}_2}{X_2} \right)^2$$

But $\mathcal{E} = \frac{q_1 \hat{\mathbf{X}}_1}{X_1} + \frac{q_2 \hat{\mathbf{X}}_2}{X_2}$ is just the electric field generated by a pair of point charges, $4\pi\epsilon_0 q_1$ at $\mathbf{X}_1$ and $4\pi\epsilon_0 q_2$ at $\mathbf{X}_2$, in two-dimensional electrostatics using SI units. The electrostatic energy, including the divergent self energies, of this configuration is exactly

$$W = \frac{q_1^2}{2\pi\epsilon_0} \ln \frac{R}{r_v} + \frac{q_2^2}{2\pi\epsilon_0} \ln \frac{R}{r_v} + \frac{q_1^2}{\pi\epsilon_0} \ln \frac{X_{12}}{r_v} \quad (3.9)$$

where $\mathbf{X}_{12}$ is the vector from vortex 1 to vortex 2. Alternatively³¹, we can express the electrostatic energy as the integral of $\frac{\epsilon_0}{2} \mathcal{E}^2$. Thus, upon comparison, the energy of the two vortex system is




Figure 3.3: Intervortex forces: top, two vortices of opposite vorticity attract; bottom, two vortices with same sense vorticity repel.

$$E_{12} = S^2 J \pi \left(q_1^2 \ln \frac{R_S}{r_v} + q_2^2 \ln \frac{R_S}{r_v} + 2q_1 q_2 \ln \frac{X_{12}}{r_v} \right) \quad (3.10)$$

Similarly, for a collection of n vortices, with cores widely separated, the spin field pattern is

$$\begin{aligned} \phi_{tot} &= \sum_{i=1}^n q_i \chi(\mathbf{X}_i) \\ \theta_{tot} &= \sum_{i=1}^n \theta_v(\mathbf{r} - \mathbf{X}_i) \approx 0 \end{aligned} \quad (3.11)$$

Following the same analogy to electrostatics as before, we find the energy of the collection of vortices is now

$$E_{tot} = S^2 J \pi \sum_{i=1}^n q_i^2 \ln \frac{R_S}{r_v} + 2S^2 J \pi \sum_{i \neq j} q_i q_j \ln \frac{X_{ij}}{r_v} \quad (3.12)$$

The force $\mathbf{F}_{ij}$ acting on vortex j due to vortex i , separated by distance X_{ij}

$$\begin{aligned}\mathbf{F}_{ij} &= -\nabla_{\mathbf{X}_{ij}} E_{tot} \\ &= \frac{S^2 J 2\pi q_i q_j}{X_{ij}} \hat{\mathbf{X}}_{ij}\end{aligned}\tag{3.13}$$

where $\hat{\mathbf{X}}_{ij}$ is a unit vector pointing from the center of vortex i to the center of vortex j . Thus, if the two vortices have the same sense, or the same sign vorticities q_i and q_j , the force is repulsive, and conversely, for opposite senses the force is attractive. Note, since in this approximation there is no interaction between the two vortex cores, the direction of the spins out of the plane at the cores—the polarization—is irrelevant.

3.2 The gyrotropic force and the vortex momentum

3.2.1 The gyrotropic force

The vortex is a stationary solution of the system. If we assume that it now moves at a small velocity $\dot{\mathbf{X}}$, for the moment with no deformation to the vortex profile, the $p\dot{q}$ action term, called the Berry's phase in a spin system, is no longer vanishing. The Berry's phase, ω_B , is a phase accumulated by the changing spin field

$$\omega_B = \int dt \int \frac{d^2 r}{a^2} S \cos \theta \dot{\phi}\tag{3.14}$$

Considering a single spin, we can interpret this phase geometrically as the solid angle swept out by the motion mapped onto the spin sphere. This is clear when we make the change of variable

$$\omega_B = S \int d\phi \cos \theta = S \int d\omega'$$

where $d\omega'$ is the area increment on the unit sphere. Refer to Figure 3.4.

We treat this term as a potential and calculate the corresponding force acting on the vortex. Let the vortex profile move as a function of $\mathbf{r} - \mathbf{X}(t)$, where $\mathbf{X}(t)$ is the center coordinate of the vortex. The Berry's phase term in the Lagrangian becomes

$$-S \int \frac{d^2 r}{a^2} \dot{\phi}_v \cos \theta_v = S \int \frac{d^2 r}{a^2} \dot{\mathbf{X}} \cdot \nabla \phi_v \cos \theta_v$$

The gyrotropic force arising from this term is found by varying it with respect to the center coordinate of the vortex^{59,65}, without the usual negative sign since




Figure 3.4: The spin path mapped onto the unit sphere. The area traced out by its motion gives the Berry's phase.

we take the term from the Lagrangian,

$$\begin{aligned}\mathbf{F}_{gyro} &= S \partial_{\mathbf{X}} \int \frac{d^2 r}{a^2} \dot{\mathbf{X}} \cdot \nabla \phi_v \cos \theta_v \\ &= -S \int \frac{d^2 r}{a^2} \nabla \left(\dot{\mathbf{X}} \cdot \nabla \phi_v \cos \theta_v \right)\end{aligned}\quad (3.15)$$

But the integrand is strictly a function of $\mathbf{r} - \mathbf{X}$ so that $\partial_{\mathbf{X}} \rightarrow -\nabla$, where ∇ is understood to be with respect to $\mathbf{r}$. Note that $\nabla^2 \phi_v = 0$ and thus

$$\nabla \left(\dot{\mathbf{X}} \cdot \nabla \phi_v \cos \theta_v \right) = \left(\dot{\mathbf{X}} \cdot \nabla \phi_v \right) \nabla \cos \theta_v$$

Using the cross-product relation $\mathbf{A} \times (\mathbf{B} \times \mathbf{C}) = (\mathbf{A} \cdot \mathbf{C}) \mathbf{B} - (\mathbf{A} \cdot \mathbf{B}) \mathbf{C}$, we find

$$-S \int \frac{d^2 r}{a^2} \left(\dot{\mathbf{X}} \cdot \nabla \phi_v \right) \nabla \cos \theta_v = S \int \frac{d^2 r}{a^2} (\nabla \cos \theta_v \times \nabla \phi_v) \times \dot{\mathbf{X}} - \left(\dot{\mathbf{X}} \cdot \nabla \cos \theta_v \right) \nabla \phi_v$$

where now both terms on the right are integrable. Consider the first term, noting that

$$(\nabla \cos \theta_v \times \nabla \phi_v)_z = \frac{\partial \cos \theta_v}{\partial x} \frac{\partial \phi_v}{\partial y} - \frac{\partial \cos \theta_v}{\partial y} \frac{\partial \phi_v}{\partial x} = \frac{\partial(\cos \theta_v, \phi_v)}{\partial(x, y)}$$

Clearly, since $\nabla \cos \theta_v$ and $\nabla \phi_v$ both lie entirely in the plane, the $\hat{\mathbf{z}}$ component is the only non-zero component. The first integral becomes

$$S \int \frac{d^2 r}{a^2} \nabla \cos \theta_v \times \nabla \phi_v = \frac{S}{a^2} \int d \cos \theta_v d \phi_v \hat{\mathbf{z}} = -\frac{2\pi S p q}{a^2} \hat{\mathbf{z}}$$




Figure 3.5: The gyrotropic force: left, a vortex with $p = 1$ and $q = -1$ traveling to the right experiences an upward force; right, a vortex with $p = 1$ and $q = 1$ traveling to the right experiences a downward force. Note $\hat{\mathbf{z}}$ is defined out of the page.

where p is the polarization of the vortex core and q is the vorticity of the vortex.

For the second integral, consider axes $x_{\parallel}$ and $x_{\perp}$ parallel and perpendicular to $\dot{\mathbf{X}}$, where the second is aligned such that $\hat{\mathbf{z}} \times \dot{\mathbf{X}} = \mathbf{x}_{\perp}$. In polar coordinates defined for this frame, the integral can be written

$$-\frac{S}{a^2} \int d^2r \left(\dot{\mathbf{X}} \cdot \nabla \cos \theta_v \right) \nabla \phi_v = -\frac{Sq}{a^2} \int dr d\chi \dot{X} \frac{d \cos \theta_v}{dr} \cos \chi (-\sin \chi, \cos \chi)$$

where we decompose $\hat{\chi} = (-\sin \chi, \cos \chi)$ into the $(x_{\parallel}, x_{\perp})$ basis. Evaluating this gives

$$-\frac{S}{a^2} \int d^2r \left(\dot{\mathbf{X}} \cdot \nabla \cos \theta_v \right) \nabla \phi_v = \pi pq \dot{X} \mathbf{x}_{\perp} = \frac{\pi Spq}{a^2} \hat{\mathbf{z}} \times \dot{\mathbf{X}} \quad (3.16)$$

The gyrotropic force is then

$$\mathbf{F}_{gyro} = -\frac{\pi Spq}{a^2} \hat{\mathbf{z}} \times \dot{\mathbf{X}} \quad (3.17)$$

Note, this result differs by a factor of 2 from that of Huber²⁵ using the formalism of Thiele, found for a magnetic domain wall⁶⁵. Thiele's starting point for the kinetic term was

$$-\cos \theta_v \dot{\phi}_v + \frac{d}{dt} \cos \theta_v \phi_v \quad (3.18)$$

which is exactly twice our starting point that includes only the first of these two terms.

Notice that the gyrotropic force is derivable from the equivalent Lagrangian term

$$\mathcal{L}_{gyro} = \frac{\pi Spq}{2a^2} \mathbf{X} \times \dot{\mathbf{X}} \cdot \hat{\mathbf{z}} = -\frac{\pi Spq}{2a^2} \mathbf{X} \times \hat{\mathbf{z}} \cdot \dot{\mathbf{X}} \quad (3.19)$$

3.2.2 The vortex momentum

The gyrotropic force can be written in the suggestive form

$$\mathbf{F}_{gyro} = -\frac{d\mathbf{P}_{gyro}}{dt} \quad (3.20)$$

where $\mathbf{P}_{gyro}$ is a momentum term from the equivalent Lagrangian term (3.19) written in the form $\mathbf{P} \cdot \dot{\mathbf{X}}$

$$\mathbf{P}_{gyro} = -\frac{\pi Spq}{2a^2} \mathbf{X} \times \hat{\mathbf{z}} \quad (3.21)$$

We now examine a direct evaluation of the vortex momentum as given in a general field theory by the operator⁵⁴

$$\mathbf{P} = -\int d^2r \tilde{\pi}(\mathbf{r}, t) \nabla \tilde{\phi}(\mathbf{r}, t) \quad (3.22)$$

where $\tilde{\pi}$ is the conjugate momentum density to the field variable $\tilde{\phi}$ (the tilde's are there to differentiate the field variable here to the azimuthal angle ϕ used previously). This operator is chosen because it is the infinitesimal generator of spatial translations, eg.

$$\begin{aligned} \tilde{\phi}(\mathbf{r} + \delta\mathbf{r}) &= \tilde{\phi}(\mathbf{r}) + \delta\tilde{\phi}(\mathbf{r}) = \tilde{\phi}(\mathbf{r}) + \nabla\tilde{\phi}(\mathbf{r}) \cdot \delta\mathbf{r} \\ \delta\tilde{\phi}(\mathbf{r}) &= \{\delta\mathbf{r} \cdot \mathbf{P}, \tilde{\phi}(\mathbf{r})\} = \nabla\tilde{\phi}(\mathbf{r}) \cdot \delta\mathbf{r} \end{aligned}$$

where we use the Poisson bracket here as defined in equation A.8 (note there we used q for $\tilde{\phi}$ and p for $\tilde{\pi}$).

For the magnetic vortex, this gives the momentum expression

$$\mathbf{P} = \int d^2r \frac{S}{a^2} \cos\theta_v \nabla\phi_v \quad (3.23)$$

Before attempting to evaluate this expression, first note that the $1/r$ behaviour in $\nabla\phi_v$ is balanced by the r in the integration measure so that the integrand is nowhere divergent.

If we blindly set the vortex at the origin of the integration region, the $\hat{\chi}_r$ direction of the integrand sums to 0 by symmetry, there being no other angular

dependence. The integral is non-zero, however, if we displace the vortex by $\mathbf{X}$ from the origin.

To evaluate this integral note that

$$\nabla \phi_v = -q\hat{\mathbf{z}} \times \nabla \ln |\mathbf{r} - \mathbf{X}| \quad (3.24)$$

Considering the momentum integral one component at a time, first the y component

$$\begin{aligned} \int d^2r \cos \theta_v \partial_x \ln |\mathbf{r} - \mathbf{X}| &= - \int dx dy \partial_x \cos \theta_v \left(\ln r - \frac{\mathbf{r} \cdot \mathbf{X}}{r^2} \right) \\ &= \int dr d\chi_r \partial_r \cos \theta_v \cos \chi_r \hat{\mathbf{r}} \cdot \mathbf{X} \\ &= -\pi q p X \end{aligned}$$

where X is the x component of $\mathbf{X}$. We expanded the $\ln$ above and truncated the series to $\mathcal{O}(1/r)$. This is in keeping with the $r \rightarrow 0$ behaviour noted in the original integral. Of course, for $r \rightarrow \infty$ the integrand decays to zero exponentially as before.

After the analogous treatment for y , we find the momentum is exactly the $\mathbf{P}_{gyro}$ describing the gyrotropic force

$$\mathbf{P}_{gyro} = -\frac{\pi S q p}{a^2} \mathbf{X} \times \hat{\mathbf{z}} \quad (3.25)$$

What does it mean exactly to have a momentum that is speed independent and coordinate dependent? Isn't this extremely bizarre? Recalling the problem of a charged particle in a magnetic field, the momentum of such a particle is modified by the presence of the magnetic field according to³⁷

$$\mathbf{p} \rightarrow \mathbf{p} - \frac{e}{c} \mathbf{A} \quad (3.26)$$

where $\mathbf{A}$ is the vector potential describing the magnetic field $\mathbf{B} = \nabla \times \mathbf{A}$, e is the electric charge and c here is the speed of light.

For the magnetic vortex, this momentum term must also correspond to a vector potential term. Completing the analogy, using $4\pi\epsilon_0 q$ as charge as in section 3.1, replacing the speed of light by the speed of magnons SJa^2/r_v , we find an effective perpendicular magnetic field $\mathbf{B} = \frac{S^2 J}{4\epsilon_0 r_v} p \hat{\mathbf{z}}$.

To further explore this interpretation, we expect the gyro-momentum to be gauge dependent. That is, we should be able to rewrite the vector potential

$$\mathbf{A} \rightarrow \mathbf{A} + \nabla_r f(r) \quad (3.27)$$

for any continuous function $f(r)$, changing the momentum expression $\mathbf{P}_{gyro}$, however, and still describe the same physical system.

Considering this gauge change in reverse, we use the gauge freedom of the Berry's phase. The Berry's phase is written in a general $\hat{\mathbf{\Omega}}$ basis

$$\omega_B = \int dt d^2r \mathbf{A}(\hat{\mathbf{\Omega}}) \dot{\hat{\mathbf{\Omega}}} \quad (3.28)$$

where $\mathbf{A}$ is a unit magnetic monopole vector potential. We change the gauge of this vector potential $\mathbf{A}$ via

$$\mathbf{A} \rightarrow \mathbf{A} + \nabla_{\hat{\mathbf{\Omega}}} f \quad (3.29)$$

where f is a general function of $\hat{\mathbf{\Omega}}$.

The momentum of the magnetic vortex is altered by noting the correspondence

$$\begin{aligned} \omega_B &= \int dt d^2r \mathbf{A}(\hat{\mathbf{\Omega}}) \dot{\hat{\mathbf{\Omega}}} \\ \mathbf{P}_{j,gyro} &= - \int d^2r A_i \nabla_{r_j} \hat{\Omega}_i \end{aligned}$$

The Berry's phase gauge change shifts the momentum definition according to

$$\mathbf{P} = \int d^2r \frac{S}{a^2} \cos \theta_v \nabla_{\mathbf{r}} \phi_v + \nabla_{\hat{\Omega}_i} f \nabla_{\mathbf{r}} \hat{\Omega}_i \quad (3.30)$$

But

$$\nabla_{\hat{\Omega}_i} f \nabla_{\mathbf{r}} \hat{\Omega}_i = \nabla_{\mathbf{r}} f(\hat{\mathbf{\Omega}}) = -\nabla_{\mathbf{X}} f(\hat{\mathbf{\Omega}})$$

since $\hat{\mathbf{\Omega}} = \hat{\mathbf{\Omega}}(\mathbf{r} - \mathbf{X})$. Thus, the additional term to the vortex momentum becomes

$$- \int d^2r \nabla_{\hat{\Omega}_i} f \nabla_{\mathbf{r}} \hat{\Omega}_i = \nabla_{\mathbf{X}} \int d^2r f = \nabla_{\mathbf{X}} F(\mathbf{X}) \quad (3.31)$$

where $F(\mathbf{X}) = \int d^2r f$ is now some general function of $\mathbf{X}$.

A continuous function $F(\mathbf{X})$ can always be expressed as the integral over another function $f(\mathbf{r}, \mathbf{X})$. Thus, the gauge freedom in the Berry's phase allows exactly the necessary gauge freedom in the gyrotropic momentum term, supporting our vector potential interpretation.




Figure 3.6: Sequenced photographs of a pair of fluid vortices with same sense vorticity. Photos were taken at 2 second intervals³⁶.

3.3 Motion of vortex pairs

Consider the motion of a pair of vortices, separated enough that the cores do not significantly interact, with polarization p_i and vorticity q_i , $i = 1, 2$. The motion so far is dictated by the balance of the inter-vortex and gyrotropic forces acting on each vortex

$$\begin{aligned} \frac{2\pi S^2 J q_1 q_2}{X_{12}^2} (\mathbf{X}_1 - \mathbf{X}_2) - \pi p_1 q_1 \hat{\mathbf{z}} \times \dot{\mathbf{X}}_1 &= 0 \\ \frac{2\pi S^2 J q_1 q_2}{X_{12}^2} (\mathbf{X}_2 - \mathbf{X}_1) - \pi p_2 q_2 \hat{\mathbf{z}} \times \dot{\mathbf{X}}_2 &= 0 \end{aligned} \quad (3.32)$$

or taking the cross-product of each equation with $\hat{\mathbf{z}}$

$$\dot{\mathbf{X}}_1 = -\frac{p_1 q_2}{p_2 q_1} \dot{\mathbf{X}}_2 = \frac{2S^2 J p_1 q_2}{X_{12}^2} \hat{\mathbf{z}} \times (\mathbf{X}_2 - \mathbf{X}_1)$$

In the case $p_1 q_2 = p_2 q_1$, $\dot{\mathbf{X}}_1 = -\dot{\mathbf{X}}_2$ and the vortices move on a common circular orbit, keeping separation X_{12} with angular frequency $\omega = \frac{4S^2 J p_1 q_2}{X_{12}^2}$ where $\omega > 0$ denotes counter-clockwise rotation.

For the opposite case, $p_1 q_2 = -p_2 q_1$, we have $\dot{\mathbf{X}}_1 = \dot{\mathbf{X}}_2$ and the vortex pair move with a common velocity on parallel lines (upward for $p_2 q_1 > 0$ and downward for $p_2 q_1 < 0$).

In this approximation, the dynamics of the vortex pair is identical to the analogous motion of a pair of fluid vortices. Referring to Figures (3.6) and (3.7), a pair of fluid vortices with the same direction circulation move in a common circular motion while a pair of opposite circulation move along parallel paths. There is the notable difference, of course, that here the type of motion is dictated by the products pq rather than just q as in regular fluid dynamics.




Figure 3.7: Sequenced photographs of a pair of fluid vortices with opposite sense vorticity. Photos were taken at 4 minute intervals³⁶.

3.4 Vortex mass

Up to now, we've assumed that the vortex profile is rigid when in motion. In fact, the profile is modified linearly in $\dot{X}$.

Assuming that the vortex moves at small velocity $\dot{X}$, expand about the rigid vortex profile

$$\begin{aligned}\phi_v &= \phi_v^{(0)}(\mathbf{r} - \mathbf{X}) + \phi_v^{(1)} \\ \theta_v &= \theta_v^{(0)}(\mathbf{r} - \mathbf{X}) + \theta_v^{(1)}\end{aligned}$$

Substituting this into the equations of motion, (3.2), to first order in $\dot{X}$, making use of the zeroth order equations of motion, these reduce to

$$\begin{aligned}-\frac{\sin \theta_v^{(0)}}{c} \dot{\mathbf{X}} \cdot \nabla \phi_v^{(0)} &= -\nabla^2 \theta_v^{(1)} - \cos 2\theta_v^{(0)} \left(\frac{1}{r_v^2} - \left(\nabla \phi_v^{(0)} \right)^2 \right) \theta_v^{(1)} \\ &\quad + \sin 2\theta_v^{(0)} \nabla \phi_v^{(1)} \cdot \nabla \phi_v^{(0)} \\ -\frac{1}{c} \dot{\mathbf{X}} \cdot \nabla \theta_v^{(0)} &= \sin \theta_v^{(0)} \nabla^2 \phi_v^{(1)} + 2 \cos \theta_v^{(0)} \left(\nabla \theta_v^{(0)} \cdot \nabla \phi_v^{(1)} + \nabla \theta_v^{(1)} \cdot \nabla \phi_v^{(0)} \right)\end{aligned}\tag{3.33}$$

Using the asymptotic expansion of the stationary vortex (3.4), the asymptotic forms of the profile perturbation become, keeping only the dominant terms in the $r \rightarrow 0$ and $r \rightarrow \infty$ limits

$$\begin{aligned}\theta_v^{(1)} &= \begin{cases} -\frac{qc_1 \dot{X} r^2}{2cr_p^2} \cos \chi, & r \rightarrow 0 \\ -\frac{q \dot{X} r^2}{cr} \cos \chi, & r \rightarrow \infty \end{cases} \\ \phi_v^{(1)} &= \begin{cases} -\frac{\dot{X} r}{2c} \sin \chi, & r \rightarrow 0 \\ -\frac{\dot{X} c_2 r_v^{3/2}}{c} \frac{e^{-r/r_v}}{\sqrt{r}} \sin \chi, & r \rightarrow \infty \end{cases}\end{aligned}\tag{3.34}$$

where c_1 and c_2 are free parameters in the unperturbed asymptotic form (3.4).

Substituting these asymptotic expressions into the energy integral, we find energy terms that are quadratic to lowest order in $\dot{X}$ (the linear terms integrate to zero by symmetry) interpretable as a $\frac{1}{2}M_v\dot{X}^2$ kinetic term:

$$E = E_v^{(0)} + E_{core}^{(1)} + E_\infty^{(1)} \quad (3.35)$$

where $E_{core}^{(1)}$ accounts for the $r = 0..r_v$ and $E_\infty^{(1)}$ accounts for the remaining $r = r_v..\infty$. Evaluated,

$$\begin{aligned} E_{core}^{(1)} &= \frac{\pi q^2 \dot{X}^2}{8Ja^4} \left(3c_1^2 r_v^2 - \frac{1}{6}c_1^2 r_v^2 - \frac{5}{12}c_1^4 r_v^2 \right) \\ E_\infty^{(1)} &= \frac{\pi \dot{X}^2}{2Ja^4} \left(q^2 r_v^2 \ln \frac{R_S}{r_v} + \frac{r_v^2}{2} + \frac{c_2^2 r_v^2}{2e^2} \right) \end{aligned} \quad (3.36)$$

Assuming an energy correction of the form

$$\Delta E = \frac{1}{2}M_v\dot{X}^2 \quad (3.37)$$

the leading term describing the vortex mass is deduced as

$$M_v = \frac{\pi q^2 r_v^2}{Ja^4} \ln \frac{R_S}{r_v}. \quad (3.38)$$

Note that this mass is, in fact, identical to the mass estimate suggested by Duan and Leggett¹¹ based on purely dimensional arguments, $M_v = \frac{E_v}{(c/r_v)^2}$.

3.5 Quantization of magnetic vortices

Quantum fluctuations in a system introduce a zero-point energy. In the previous chapter, we quantized the magnons finding this zero-point energy to be $\sum \frac{1}{2}\hbar\omega_k$, summed over the entire k -spectrum. In the presence of a magnetic vortex, the magnon spectrum is shifted. Since we prefer to have a consistent definition of the magnons and vortices, the shift in the zero-point energy of the magnons is associated with the quantized vortex.

Quantization of a magnetic vortex involves quantizing the small variations about it and examining how the energy of these modes shift from the analogous modes in the absence of a vortex⁵⁰. See Appendix B.1 for more details.

Expanding θ and ϕ about a vortex, $\theta = \theta_v + \vartheta$ and $\phi = \phi_v + \varphi$, in the non-linear equations of motion (3.2), yields the linearized equations in ϑ and φ

$$\begin{aligned}
\frac{\sin \theta_v}{c} \frac{\partial \varphi}{\partial t} &= -\nabla^2 \vartheta - \cos 2\theta_v \left(\frac{1}{r_v^2} - (\nabla \phi_v)^2 \right) \vartheta + \sin 2\theta_v \nabla \varphi \cdot \nabla \phi_v \\
\frac{1}{c} \frac{\partial \vartheta}{\partial t} &= \sin \theta_v \nabla^2 \varphi + 2 \cos \theta_v (\nabla \theta_v \cdot \nabla \varphi + \nabla \vartheta \cdot \nabla \phi_v)
\end{aligned} \tag{3.39}$$

These are very similar to the vacuum magnon equations of motion with the addition of a few perturbing terms. Notably, these additional terms all decay away the vortex core and will be treated in a Born approximation (applicability of this approximation is discussed at the end of the next section).

Alternatively^{28,29,56}, we could expand as

$$\begin{aligned}
\theta &= \theta_v + \vartheta \\
\phi &= \phi_v + \frac{\varphi}{\sin \theta_v}
\end{aligned} \tag{3.40}$$

yielding the linearized equations in ϑ and φ

$$\begin{aligned}
\frac{1}{c} \frac{\partial \varphi}{\partial t} &= -\nabla^2 \vartheta - \cos 2\theta_v \left(\frac{1}{r_v^2} - (\nabla \phi_v)^2 \right) \vartheta + 2 \cos \theta_v \nabla \varphi \cdot \nabla \phi_v \\
\frac{1}{c} \frac{\partial \vartheta}{\partial t} &= \nabla^2 \varphi + \cos^2 \theta_v \left(\frac{1}{r_v^2} - (\nabla \phi_v)^2 \right) \varphi + (\nabla \theta_v)^2 \varphi + 2 \cos \theta_v \nabla \vartheta \cdot \nabla \phi_v
\end{aligned}$$

or, equivalently, in the more symmetric form

$$\begin{aligned}
\frac{r_v}{c} \frac{\partial \varphi}{\partial t} &= (-\nabla^2 + V_1(x)) \vartheta + \frac{2q \cos \theta_v}{x^2} \frac{\partial \varphi}{\partial \xi} \\
-\frac{r_v}{c} \frac{\partial \vartheta}{\partial t} &= (-\nabla^2 + V_2(x)) \varphi - \frac{2q \cos \theta_v}{x^2} \frac{\partial \vartheta}{\partial \xi}
\end{aligned} \tag{3.41}$$

where $x = \frac{r}{r_v}$, radial derivatives are now with respect to x , and

$$\begin{aligned}
V_1(x) &= \left((\nabla \phi_v)^2 - 1 \right) \cos 2\theta_v \\
V_2(x) &= \left((\nabla \phi_v)^2 - 1 \right) \cos^2 \theta_v - (\nabla \theta_v)^2
\end{aligned} \tag{3.42}$$

This form is particularly suitable for examining the core effects and searching for possible bound modes.

3.5.1 Phase shifts in the Born approximation

The perturbing terms are localized to the vortex and can be treated in a Born approximation. The dominant scattering term decays as $\frac{1}{r^2}$, whereas the remaining terms, neglected in the following treatment, die exponentially. The error introduced by neglecting these terms will be discussed in the final analysis at the end of this section. The magnon equations of motion are modified to

$$\begin{aligned}\frac{1}{c} \frac{\partial \varphi}{\partial t} &= -\nabla^2 \vartheta + \left(\frac{1}{r_v^2} - \frac{q^2}{r^2} \right) \vartheta \\ \frac{1}{c} \frac{\partial \vartheta}{\partial t} &= \nabla^2 \varphi\end{aligned}\tag{3.43}$$

The perturbation treatment is most straightforward in a single variable. Eliminating the ϑ variable, we have

$$\frac{1}{c^2} \frac{\partial^2 \varphi}{\partial^2 t} = -\nabla^4 \varphi + \left(\frac{1}{r_v^2} - \frac{q^2}{r^2} \right) \nabla^2 \varphi\tag{3.44}$$

Note the additional term $\frac{q^2}{r^2}$ modifying the vacuum equations of motion of the magnons (2.3). The Born approximation is applied using the standard partial-wave analysis from scattering theory³⁸.

Consider the orthonormal basis functions $\xi_{\mathbf{k}}$ such that $\nabla^2 \xi_{\mathbf{k}} \rightarrow -k^2 \xi_{\mathbf{k}}$ and assume harmonic time dependence. We expand φ in this basis

$$\varphi = \sum_{\mathbf{k}'} c_{\mathbf{k}'} e^{i\omega_{\mathbf{k}'} t} \xi_{\mathbf{k}'}\tag{3.45}$$

where to zeroth order we've assumed

$$c_{\mathbf{k}'}^{(0)} = \begin{cases} 1, & \mathbf{k}' = \mathbf{k}; \\ 0, & \text{otherwise.} \end{cases}\tag{3.46}$$

The zeroth order terms simply reduce to the vacuum equations of motion. The first order terms are

$$-\sum_{\mathbf{k}' \neq \mathbf{k}} \frac{\omega_{\mathbf{k}'}^2}{c^2} c_{\mathbf{k}'}^{(1)} \xi_{\mathbf{k}'} = -\sum_{\mathbf{k}' \neq \mathbf{k}} k'^2 Q'^2 c_{\mathbf{k}'}^{(1)} \xi_{\mathbf{k}'} + \frac{q^2 k^2}{r^2} \xi_{\mathbf{k}}\tag{3.47}$$

where we've cancelled the common $e^{i\omega_{\mathbf{k}} t}$ factor. Recall $Q^2 = k^2 + \frac{1}{r_v^2}$. Multiplying by $\xi_{\mathbf{k}'}^\dagger$, and integrating over space, enforcing orthonormality of the $\{\xi_{\mathbf{k}}\}$, we find an expression for the first order coefficients

$$c_{\mathbf{k}'}^{(1)} = -\frac{c^2}{\omega_{\mathbf{k}}^2 - c^2 k'^2 Q'^2} \int d^2 r \xi_{\mathbf{k}'}^\dagger \frac{q^2 k^2}{r^2} \xi_{\mathbf{k}}\tag{3.48}$$

Substituting for the unperturbed magnon spectrum (2.4) and using plane waves for the orthonormal basis, the first order correction to φ is

$$\varphi^{(1)}(\mathbf{r}) = - \int \frac{d^2 k'}{(2\pi)^2} \frac{e^{i\mathbf{k}' \cdot \mathbf{r}}}{k^2 Q^2 - k'^2 Q'^2} \int d^2 r' e^{-i\mathbf{k}' \cdot \mathbf{r}'} \frac{q^2 k^2}{r'^2} e^{i\mathbf{k} \cdot \mathbf{r}'}$$

First, integrating over the polar angle $\phi_{\mathbf{k}'}$ from 0 to π , we obtain

$$\varphi^{(1)}(\mathbf{r}) = \frac{1}{4} \int_{-\infty}^{\infty} \frac{dk' k'}{2\pi} \frac{H_0^{(1)}(k|\mathbf{r} - \mathbf{r}'|) + H_0^{(2)}(k|\mathbf{r} - \mathbf{r}'|)}{k'^2 Q'^2 - k^2 Q^2 \pm i\epsilon} \int d^2 r' e^{-i\mathbf{k}' \cdot \mathbf{r}'} \frac{q^2 k^2}{r'^2} e^{i\mathbf{k} \cdot \mathbf{r}'}$$

The $\pm i\epsilon$ are chosen to displace the poles so as to pick the outgoing wave (the plus is for the $H_0^{(2)}$ integral, the minus for the $H_0^{(1)}$ integral). Considering the $H_0^{(2)}$ integral, there are poles in the complex k' plane at $k' = \pm(k + i\epsilon')$. Noting the asymptotic behaviour

$$H_0^{(2)}(kr) \rightarrow \sqrt{\frac{2}{\pi kr}} e^{-i(kr - (l + \frac{1}{2})\frac{\pi}{2})} \quad (3.49)$$

we close the contour about the positive imaginary axis for the $H_0^{(2)}$ integral, to pick up the $k' = k + i\epsilon$ pole with residue $\frac{1}{2(k^2 + Q^2)} H_0^{(2)}(k|\mathbf{r} - \mathbf{r}'|)$. The integral over $H_0^{(1)}$ is just the complex conjugate (c.c) of that over $H_0^{(2)}$ and hence follows immediately.

Thus we have

$$\varphi^{(1)}(\mathbf{r}) = \frac{ik^2}{8(k^2 + Q^2)} \int d^2 r' H_0^{(2)}(k|\mathbf{r} - \mathbf{r}'|) \frac{q^2}{r'^2} e^{i\mathbf{k} \cdot \mathbf{r}'} + c.c.$$

We now expand the Hankel functions according to the identity

$$H_0^{(1,2)}(k|\mathbf{r} - \mathbf{r}'|) = \sum_{l=\infty}^{\infty} J_l(kr') H_l^{(1,2)}(kr) e^{il(\phi - \phi')} \quad (3.50)$$

if $r > r'$, an allowable assumption if we only want the wavefunction correction for asymptotic r , and ϕ (ϕ') is the polar angle of $\mathbf{r}$ ($\mathbf{r}'$). Expand the plane wave as

$$e^{i\mathbf{k} \cdot \mathbf{r}'} = \sum_{m=-\infty}^{\infty} i^m J_m(kr') e^{im\phi'} \quad (3.51)$$

After integration over ϕ' (giving a factor $2\pi\delta_{lm}$), the wavefunction correction is

$$\varphi^{(1)}(\mathbf{r}) = \frac{\pi ik^2}{4(k^2 + Q^2)} \sum_{l=\infty}^{\infty} i^l H_l^{(2)}(kr) e^{il\phi} \int dr' \frac{q^2}{r'} J_l^2(kr') + c.c. \quad (3.52)$$

Recall we assumed the unperturbed $\varphi^{(0)}$ solution was a plane wave, expandable according to (3.51), so that the entire solution can be written, up to first order,

$$\varphi(\mathbf{r}) = \sum_{l=-\infty}^{\infty} i^l e^{il\phi} \left[J_0(kr) + \left(\frac{\pi i k^2}{4(k^2 + Q^2)} H_l^{(2)}(kr) \int dr' \frac{q^2}{r'} J_l^2(kr') + c.c. \right) \right]$$

Comparing this with the sum of an incoming and outgoing wave

$$\begin{aligned} \frac{1}{2} \left(e^{i\Delta_l} H_l^{(1)}(kr) + e^{-i\Delta_l} H_l^{(2)}(kr) \right) &= J_l(kr) - i \frac{\Delta_l}{2} H_l^{(2)}(kr) + i \frac{\Delta_l}{2} H_l^{(1)}(kr) \\ &\rightarrow \sqrt{\frac{2}{\pi kr}} e^{-i\Delta_l} \cos \left(kr - \left(l + \frac{1}{2} \right) \frac{\pi}{2} - \Delta_l \right) \text{ as } r \rightarrow \infty \end{aligned} \quad (3.53)$$

gives for the phase shift of the l^{th} order wave

$$\Delta_l = -\frac{\pi}{2} \frac{k^2}{k^2 + Q^2} \int dr' \frac{q^2}{r'} J_l^2(kr') \quad (3.54)$$

These phase shifts perturb the magnon wavevector $k = k^{scatt} - \Delta_l$, and, hence, the magnon spectrum ω_k . For proper counting of the total energy shift, first discretize k by fixing the boundary conditions of the wavefunction at $r = R_S$ so that

$$\pi n = k_n R_S = k_n^{scatt} R_S - \Delta_l(k_n) \quad (3.55)$$

Notice that asymptotically we have a cosine wavefunction as opposed to a plane wave as described by Rajaraman⁵⁰. Letting the system size tend to infinity then

$$\sum_{k_n} \rightarrow \frac{R_S}{\pi} \int dk$$

The zero point energy shift, given by the change in the zero point energy of the small oscillation modes when the vortex is present as compared to those in vacuum, is then

$$\Delta E = \frac{1}{2} \sum_{k,l} \hbar \delta \omega_k \quad (3.56)$$

But $k = k^{scatt} - \Delta_l$,

$$\begin{aligned}
\delta\omega_k &= \omega(k^{scatt}) - \omega(k) \\
&= \omega\left(k + \frac{\Delta_l}{R_S}\right) - \omega(k) \\
&= \frac{\partial\omega(k)}{\partial k} \frac{\Delta_l(k)}{R_S}
\end{aligned}$$

so that

$$\Delta E = \frac{\hbar}{2} \sum_{l=-\infty}^{\infty} \int \frac{dk}{\pi} \frac{\partial\omega}{\partial k} \Delta_l(k) \quad (3.57)$$

Substituting for $\Delta_l(k)$ from equation (3.54), noting that $\frac{\partial\omega}{\partial k} = c \frac{k^2 + Q^2}{Q}$,

$$\begin{aligned}
\Delta E &= -\frac{\hbar c}{4} \int dk \frac{k^2}{k^2 + Q^2} \frac{k^2 + Q^2}{Q} \int dr' \frac{q^2}{r'} \sum_{l=-\infty}^{\infty} J_l^2(kr') \\
&= -\frac{\hbar c q^2}{4} \int dk \frac{k^2}{Q} \ln \frac{R_S}{r_v}
\end{aligned}$$

where we've used that $\sum_{l=-\infty}^{\infty} J_l^2(kr') = 1$. Note that the radial integral is cut off by the vortex core size. This is because the perturbing term changes behaviour drastically in the core so that our analysis cannot be extended there. The k integral can be evaluated noting that

$$\frac{d}{dk} \left(\frac{1}{2} k Q - \frac{1}{2r_v^2} \ln(k + Q) \right) = \frac{k^2}{Q}$$

The result is ultraviolet divergent so that we must impose a cutoff of $1/a$, physically reasonable if we recall that a is the lattice spacing of the discrete lattice. Finally, the energy shift in the presence of a magnetic vortex is

$$\Delta E = -\frac{\hbar c q^2}{4} \ln \frac{R_S}{r_v} \left(\frac{\sqrt{r_v^2 + a^2}}{2a^2 r_v} - \frac{1}{2r_v^2} \ln \frac{r_v + \sqrt{r_v^2 + a^2}}{a} \right) \quad (3.58)$$

This zero-point energy shift, due to the presence of the vortex, is associated to the quantized vortex rather than the magnons⁵⁰. Thus, ΔE is the zero-point energy of the vortex. Note that the interaction actually decreases the quantum energy of the vortex-magnon system.

We can examine the error in neglecting the exponentially decaying terms by replacing the r_v/r' behaviour by $\exp(-r/r_v)$. Essentially, this would replace the log divergence in the final result with unity. Hence, in comparison with the main contribution, these exponentially decaying terms are negligible.

The Born approximation amounts to the substitution of

$$\langle \phi_f | U | \psi_i \rangle \rightarrow \langle \phi_f | U | \phi_i \rangle \quad (3.59)$$

where ϕ and ψ denote the unperturbed and modified waveforms, respectively. In general, the validity of the Born approximation depends on how much the waveforms differ in the region of the scattering potential⁴⁶. In our case, the Born approximation indicates that the two wavefunctions in fact differ to first order by equation (3.52) which is proportional to the predicted phase shifts. This is circular reasoning; however, in the case of those quasiparticle modes delocalized over the system, we expect the waveform not to change significantly. On the other hand, there are quasiparticles that become trapped by the vortex center. Clearly, for these modes, the wavefunctions are drastically modified in the vortex core, where the scattering potential is greatest, so that a Born approximation is invalid. We examine these bound modes in the next section to show how they result from the translational symmetry broken by the vortex solution.

3.5.2 Bound modes

As pointed out by Ivanov et. al.^{28,29,56}, the short range interactions neglected in (3.43) can drastically alter the behaviour of certain modes. The symmetric perturbing equations, (3.41), are more suitable for exploring the core region. Assume a solution of the form

$$\begin{aligned} \vartheta &= f(x) \cos(m\chi + \omega t + \psi) \\ \varphi &= g(x) \sin(m\chi + \omega t + \psi) \end{aligned} \quad (3.60)$$

Substituting this into (3.41) yields equations for f and g

$$\begin{aligned} \left(\frac{d^2}{dx^2} + \frac{1}{x} \frac{d}{dx} - \frac{m^2}{x^2} - V_1(x) \right) f &= \left(\frac{wr_v}{c} + \frac{2qm \cos \theta_v}{x^2} \right) g \\ \left(\frac{d^2}{dx^2} + \frac{1}{x} \frac{d}{dx} - \frac{m^2}{x^2} - V_2(x) \right) g &= \left(\frac{wr_v}{c} + \frac{2qm \cos \theta_v}{x^2} \right) f \end{aligned} \quad (3.61)$$

recalling that $x = \frac{r}{r_v}$ and that $V_1(x)$ and $V_2(x)$ are defined in (3.42).

For $\omega = 0$, there exist exact solutions for $m = 0, \pm 1$

$$\begin{aligned} f &= m\theta'_v \\ g &= -\frac{q \sin \theta_v}{x^{|m|}} \end{aligned} \quad (3.62)$$

For $|m| > 1$, the asymptotic behaviour of the modes is entirely unbounded so that the vortex center has not greatly shifted the magnon wavefunctions and the Born approximation applied in the previous section should be valid.

Consider first the $m = 0$ result. Combining the unperturbed vortex profile with this result (recall the normalization of the perturbations as in (3.40))

$$\begin{aligned}\phi &= q\chi - q\delta\chi \\ \theta &= \theta_v\end{aligned}\tag{3.63}$$

where $\delta\chi$ is the coefficient of the linearized solution. We find that it corresponds simply to the freedom of uniform rotation in the xy -plane.

Similarly, consider the $m = \pm 1$ solutions

$$\begin{aligned}\phi &= q\chi - \frac{q\delta r}{r} \sin(m\chi - \psi) \\ \theta &= \theta_v + m\delta r \theta'_v \cos(m\chi - \psi)\end{aligned}$$

with δr as the coefficient of the linearized $m = \pm 1$ solution. But note that the additional contributions can be re-expressed as

$$\begin{aligned}\phi &= q\chi + \nabla\phi_v \cdot m\delta\mathbf{r} \\ \theta &= \theta_v + \nabla\theta_v \cdot m\delta\mathbf{r}\end{aligned}\tag{3.64}$$

where $\delta\mathbf{r}$ is now a vector of magnitude δr in the direction defined by the polar angle ψ (see Figure 3.8). Thus, these two modes represent infinitesimal motion along $\pm\delta\mathbf{r}$ (the sign chosen by the sign of m).

Clearly, these bound modes are inadequately treated using the Born approximation and must be treated separately somehow. Ivanov et. al.^{28,29,56} attempted to calculate the phase shifts of these modes separately and to subsequently use them to describe the angular and translational motion of the magnetic vortices. Alternatively, however, one can treat the problem using collective coordinates (see Appendix B.1.2 for more details) conveniently separating these so-called zero modes and treating the remaining modes in a Born approximation.

In the next chapter, we expand the interactions of the magnetic vortex with the environment magnons using collective coordinates. Using a path integral formalism, we separate the degrees of freedom of the vortex motion from those of the environment and proceed to integrate out these modes yielding the effective dynamics of the vortex.




Figure 3.8: The directions relevant to a small translation of the vortex along $\delta\mathbf{r}$.

Chapter 4

Vortex dynamics

We now have all the background to interact the vortices and magnons. Using a variety of techniques, we examine the effects of couplings between the two systems to the vortex energy and dynamics. In the previous chapter, we already saw how a modification in the magnon spectrum can be interpreted as a quantum energy shift associated with the vortex.

First, using regular perturbation theory, we examine the one magnon coupling with the vortex velocity giving rise to an inertial mass and a dissipation rate of a moving vortex. We also examine the long range two magnon coupling in this language, finding almost immediately the zero point energy shift that in the previous chapter required calculating all magnon phase shifts.

The effective vortex dynamics are derived by finding the time evolution of the vortex-magnon density matrix and tracing over the magnon modes. We use the Feynman-Vernon formalism, describing the density matrix with path integrals. We again deduce the vortex inertial mass, in agreement with perturbation results. The vortex motion is again dissipative; however, we find the vortex damping forces explicitly and characterize the associated fluctuating forces. Generalization to a collection of vortices is carried out.

In addition to the previously derived gyrotropic and inter-vortex forces, we derive microscopically vortex damping forces. We introduce for the first time in such a magnetic system a transverse damping force, analogous to the Iordanskii force acting on a vortex in a superfluid. These are derived from the action terms found in the vortex density matrix propagator (4.91) and have corresponding fluctuating forces with correlations given by (4.92).

Alternatively, we consider decomposing the motion in a Bessel function basis, $\{J_m(kX(t))e^{im\phi_x}\}$, to obtain Brownian motion for the components with an effective action given by (4.96) and corresponding fluctuating force correlations (4.98).

4.1 Vortex-magnon interaction terms

We work with the complete non-linear Lagrangian for our magnetic system

$$\mathcal{L} = S \int \frac{d^2r}{a^2} \left(-\dot{\phi} \cos \theta - \frac{c}{2} \left((\nabla \theta)^2 + \sin^2 \theta \left((\nabla \phi)^2 - \frac{1}{r_v^2} \right) \right) \right) \quad (4.1)$$

Expanding the Lagrangian density about the vortex profile via $\theta = \theta_v + \vartheta$ and $\phi = \phi_v + \varphi$ we find the following terms in the integrand

$$\begin{aligned} & \frac{S}{a^2} \left(\dot{\phi}_v + \dot{\varphi} \right) \left(-\cos \theta_v + \sin \theta_v \vartheta \right) - \frac{S^2 J}{2} \left((\vec{\nabla} \theta_v)^2 + 2 \vec{\nabla} \theta_v \cdot \vec{\nabla} \vartheta + (\vec{\nabla} \vartheta)^2 + \right. \\ & \left. (\sin^2 \theta_v + \sin 2\theta_v \vartheta + \cos 2\theta_v \vartheta^2) \left((\vec{\nabla} \phi_v)^2 - \frac{1}{r_v^2} + 2 \vec{\nabla} \phi_v \cdot \vec{\nabla} \varphi + (\vec{\nabla} \varphi)^2 \right) \right) \end{aligned}$$

The zeroth order terms in φ and ϑ simply give the vortex action; the first order terms give ϑ multiplied by the $\frac{\partial \phi_v}{\partial t}$ equation of motion and φ multiplied by the $\frac{\partial \theta_v}{\partial t}$ equation of motion and thus are zero, except, notably, the one magnon dynamic term

$$\frac{S}{a^2} \dot{\phi}_v \sin \theta_v \vartheta \quad (4.2)$$

Finally, the remaining two magnon terms are

$$\begin{aligned} & \frac{S}{a^2} \dot{\varphi} \sin \theta_v \vartheta - \frac{S^2 J}{2} \left((\vec{\nabla} \vartheta)^2 + \sin^2 \theta_v (\vec{\nabla} \varphi)^2 + 2 \sin 2\theta_v \vec{\nabla} \phi_v \cdot \vec{\nabla} \varphi \vartheta \right. \\ & \left. + \cos 2\theta_v \left((\vec{\nabla} \phi_v)^2 - \frac{1}{r_v^2} \right) \vartheta^2 \right) \quad (4.3) \end{aligned}$$

Minimizing these action terms, we find the perturbed equations of motion similar to (3.39)

$$\begin{aligned} \frac{\sin \theta_v}{c} \frac{\partial \varphi}{\partial t} + \frac{\sin \theta_v}{c} \dot{\phi}_v &= -\nabla^2 \vartheta - \cos 2\theta_v \left(\frac{1}{r_v^2} - (\nabla \phi_v)^2 \right) \vartheta + \sin 2\theta_v \nabla \varphi \cdot \nabla \phi_v \\ \frac{1}{c} \frac{\partial \vartheta}{\partial t} &= \sin \theta_v \nabla^2 \varphi + 2 \cos \theta_v (\nabla \theta_v \cdot \nabla \varphi + \nabla \vartheta \cdot \nabla \phi_v) \quad (4.4) \end{aligned}$$

Define the vortex profile relative to the center coordinate[†] $\mathbf{X}$

[†]There is no need to add a collective coordinate reflecting the rotational symmetry of the problem since this is actually just a restatement of the 2-dimensional translational freedom, already entirely taken care of in the 2-dimensional center coordinate.

$$\begin{aligned}\phi_v &= q\chi(\mathbf{r} - \mathbf{X}) \\ \theta_v &= \theta_v(\mathbf{r} - \mathbf{X})\end{aligned}\tag{4.5}$$

The center coordinates play the role of the collective coordinates in this system, introduced to account for the continuous translational symmetry broken by the vortex. They are elevated to operators.

Focussing on the one magnon perturbative term, (4.2), expanding in terms of the collective coordinates, we find

$$\frac{S}{a^2} \dot{\phi}_v \sin \theta_v \vartheta = -\frac{S}{a^2} \dot{\mathbf{X}} \cdot \nabla \phi_v \sin \theta_v \vartheta \tag{4.6}$$

In the previous chapter, this term perturbed the vortex profile under motion, introducing a finite vortex mass.

4.2 Perturbation theory results

4.2.1 Vortex mass revisited

Consider the one magnon coupling (4.6). This term can be considered as a perturbing term of the vortex profile under motion or, alternatively, as a vortex-magnon coupling. Fourier transforming ϑ according to (2.5), now with $\vartheta = \vartheta(\mathbf{r} - \mathbf{X})$, we can rewrite the coupling as

$$\begin{aligned}-\int \frac{d^2 k a^2}{(2\pi)^2} S \int \frac{d^2 r}{a^2} \sin \theta_v \dot{\mathbf{X}} \cdot \nabla \phi_v e^{-i\mathbf{k} \cdot \mathbf{r}} \vartheta_{\mathbf{k}} = \\ -\frac{S}{a^2} \int \frac{d^2 k a^2}{(2\pi)^2} e^{-i\mathbf{k} \cdot \mathbf{X}} \vartheta_{\mathbf{k}} \int d^2 r e^{-ikr \cos \chi_{kr}} \sin \theta_v \frac{q \dot{X} \sin \chi_{vr}}{r}\end{aligned}$$

where the $\mathbf{r}$ integration has been shifted to move the vortex center coordinates into the exponential. Expanding $\chi_{vr} = \chi_{vk} + \chi_{kr}$ and noting that

$$\int d\chi_{kr} \sin \chi_{kr} e^{-ikr \cos \chi_{kr}} = 0$$

the coupling term becomes

$$\begin{aligned}
& -\frac{S}{a^2} \int \frac{d^2 k a^2}{(2\pi)^2} e^{-i\mathbf{k} \cdot \mathbf{X}} \vartheta_{\mathbf{k}} \int d^2 r e^{-i k r \cos \chi_{kr}} \sin \theta_v \frac{q \dot{X} \sin \chi_{vk} \cos \chi_{kr}}{r} \\
& = \frac{2\pi i S q}{a^2} \int \frac{d^2 k a^2}{(2\pi)^2} e^{-i\mathbf{k} \cdot \mathbf{X}} \vartheta_{\mathbf{k}} \int dr \dot{X} \sin \chi_{vk} J_1(kr) \sin \theta_v \\
& = \frac{2\pi i S q}{a^2} \int \frac{d^2 k a^2}{(2\pi)^2} e^{-i\mathbf{k} \cdot \mathbf{X}} \vartheta_{\mathbf{k}} \frac{\dot{\mathbf{X}} \cdot \hat{\chi}_k}{k Q r_v}
\end{aligned}$$

where we approximate $\sin \theta_v \approx 1 - e^{-r/r_v}$, which has the right asymptotic behaviour for $r \rightarrow 0$ and $r \rightarrow \infty$. Expressing $\vartheta_{\mathbf{k}}$ in terms of creation and annihilation operators given in (2.8), we finally obtain

$$\int \frac{d^2 k a^2}{(2\pi)^2} e^{-i\mathbf{k} \cdot \mathbf{X}} \frac{2\pi q}{a^2} \sqrt{\frac{\hbar S}{2kQ}} \frac{\dot{\mathbf{X}} \cdot \hat{\chi}_k}{Q r_v} (a_{\mathbf{k}}^\dagger - a_{-\mathbf{k}}) \quad (4.7)$$

or expressed a little more symmetrically

$$\int \frac{d^2 k a^2}{(2\pi)^2} \frac{2\pi q}{a^2} \sqrt{\frac{\hbar S}{2kQ}} \frac{\dot{\mathbf{X}} \cdot \hat{\chi}_k}{Q r_v} (e^{-i\mathbf{k} \cdot \mathbf{X}} a_{\mathbf{k}}^\dagger - e^{i\mathbf{k} \cdot \mathbf{X}} a_{-\mathbf{k}}) \quad (4.8)$$

Note the similarity of the coupling here to that of the polaron problem discussed, for example, by Feynman¹⁵.

In first order perturbation theory, this coupling provides no energy shift since it necessarily changes the number of magnons between the initial and final state.

In second order perturbation theory, we consider the diagram shown in Figure 4.1 corresponding to the emission and re-absorption of a virtual magnon. The energy shift provided by this diagram, which with foresight we call E_{mass} ,

$$E_{mass} = \int \frac{d^2 k a^2}{(2\pi)^2} \frac{2\pi^2 \hbar S q^2}{a^4 k Q^3 r_v^2} \left(\dot{\mathbf{X}} \cdot \hat{\chi}_k \right)^2 \frac{1}{(\mathbf{P} - \hbar \mathbf{k})^2 / 2M_v + \hbar c k Q - P^2 / 2M_v}$$

However, so far the vortex has no inertial energy, $P^2 / 2M \rightarrow 0$, and

$$E_{mass} = \int \frac{d^2 k a^2}{(2\pi)^2} \frac{1}{\hbar c k Q} \frac{2\pi^2 \hbar S q^2}{a^4 k Q^3 r_v^2} \left(\dot{\mathbf{X}} \cdot \hat{\chi}_k \right)^2 \quad (4.9)$$

The integration over the polar angle of $\mathbf{k}$ contributes a factor π . We expand the k and Q dependence in partial fractions

$$\frac{1}{k Q^4 r_v^2} = \frac{r_v^2}{k} - \frac{k r_v^2}{Q^2} - \frac{k}{Q^4}$$

The radial integral is evaluated as




Figure 4.1: Lowest order contributing diagram for the first order vortex-magnon coupling term.

$$\begin{aligned}
 E_{mass} &= \frac{\pi q^2 \dot{X}^2}{2Ja^4} \int dk \frac{k}{k^2 Q^3 r_v^2} \\
 &= \frac{\pi q^2 \dot{X}^2}{2Ja^4} \int dk \left(\frac{r_v^2}{k} - \frac{k r_v^2}{Q^2} - \frac{k}{Q^4} \right) \\
 &= \frac{\pi q^2 r_v^2 \dot{X}^2}{2Ja^4} \left(\ln \frac{R_S}{a} - \frac{1}{2} \ln \frac{a^2 + r_v^2}{a^2} - \frac{1}{2} \frac{r_v^2}{a^2 + r_v^2} \right) \quad (4.10)
 \end{aligned}$$

where we've imposed both an upper and lower cutoff, with a the lattice spacing and R_S the system size.

Thus, identifying this as a $\frac{1}{2} M_v \dot{X}^2$ inertial term, we find a vortex mass of

$$M_v = \frac{\pi q^2 r_v^2}{Ja^4} \left(\ln \frac{R_S}{\sqrt{a^2 + r_v^2}} - \frac{1}{2} \frac{r_v^2}{a^2 + r_v^2} \right) \quad (4.11)$$

in agreement to leading order with the analysis of section 3.4. The $r_v \rightarrow \sqrt{a^2 + r_v^2}$ replacement corrects the $r_v \rightarrow 0$ limiting behaviour.

Mass tensor of a collection of vortices

We can easily generalize this result to a collection of vortices in this formalism. Recall that the n -vortex superposed solution is given by

$$\begin{aligned}
 \phi_{tot} &= \sum_{i=1}^n q_i \chi(X_i) \\
 \theta_{tot} &= \sum_{i=1}^n \theta_v(\mathbf{r} - \mathbf{X}_i) \quad (4.12)
 \end{aligned}$$

so that the one magnon coupling becomes

$$\sum_{i=1}^n \int \frac{d^2 k a^2}{(2\pi)^2} e^{-i\mathbf{k} \cdot \mathbf{X}_i} \frac{2\pi q_i}{a^2} \sqrt{\frac{\hbar S}{2kQ}} \frac{\dot{\mathbf{X}}_i \cdot \hat{\chi}_k}{Q r_v} (a_{\mathbf{k}}^\dagger - a_{-\mathbf{k}}) \quad (4.13)$$




Figure 4.2: Definition of angles for evaluation of off-diagonal mass terms.

The second order energy correction is now

$$E_{mass} = \sum_{i,j=1}^n \int d^2k \frac{q_i q_j}{2Ja^4 k^2 Q^4 r_v^2} \left(\dot{\mathbf{X}}_i \cdot \hat{\chi}_k e^{-i\mathbf{k} \cdot \mathbf{X}_i} \right) \cdot \left(\dot{\mathbf{X}}_j \cdot \hat{\chi}_k e^{i\mathbf{k} \cdot \mathbf{X}_j} \right) \quad (4.14)$$

The diagonal terms evaluate exactly as above. The off-diagonal terms can be evaluated noting that

$$\begin{aligned} & \int d\chi_{kr_{ij}} \sin \chi_{kv_1} \sin \chi_{kv_2} \exp(ikr_{ij} \cos \chi_{kr_{ij}}) \\ &= \frac{1}{2} \int d\chi_{kr_{ij}} (\cos(\chi_{kv_1} - \chi_{kv_2}) + \cos(\chi_{kv_1} + \chi_{kv_2})) \exp(ikr_{ij} \cos \chi_{kr_{ij}}) \end{aligned}$$

where $r_{ij} = |\mathbf{X}_i - \mathbf{X}_j|$. If we assume the various angles are defined as in Figure 4.2, then $\chi_{kv_1} - \chi_{kv_2} = \chi_{v_1 v_2}$ and $\chi_{kv_1} + \chi_{kv_2} = \chi_{v_1 r_{ij}} + \chi_{v_2 r_{ij}} + 2\chi_{kr_{ij}}$.

Noting by symmetry that the $\sin \chi_{kr_{ij}}$ terms integrate to zero, we have

$$\begin{aligned} & \int d\chi_{kr_{ij}} \sin \chi_{kv_1} \sin \chi_{kv_2} \exp(ikr_{ij} \cos \chi_{kr_{ij}}) \\ &= \frac{1}{2} \int d\chi_{kr_{ij}} (\cos \chi_{v_1 v_2} + \cos(\chi_{v_1 r_{ij}} + \chi_{v_2 r_{ij}}) \cos 2\chi_{kr_{ij}}) \exp(ikr_{ij} \cos \chi_{kr_{ij}}) \\ &= \pi \cos \chi_{v_1 v_2} J_0(kr_{ij}) + \pi \cos(\chi_{v_1 r_{ij}} + \chi_{v_2 r_{ij}}) J_2(kr_{ij}) \end{aligned}$$

Next we perform the integrals over k , noting that

$$\int_{1/R_S}^{1/a} dk \frac{J_0(kr_{ij})}{k} = \ln \frac{R_S}{r_{ij}}$$

$$\int_0^\infty dk \frac{J_2(kr_{ij})}{k} = \frac{1}{2}$$

and rewrite

$$\cos(\chi_{v_1 r_{ij}} + \chi_{v_2 r_{ij}}) = (\hat{\mathbf{X}}_i \cdot \hat{\mathbf{e}}_\Delta)(\hat{\mathbf{X}}_j \cdot \hat{\mathbf{e}}_\Delta) - (\hat{\mathbf{X}}_i \times \hat{\mathbf{e}}_\Delta) \cdot (\hat{\mathbf{X}}_j \times \hat{\mathbf{e}}_\Delta)$$

Finally the energy correction term becomes

$$E_{mass} = \sum_{i,j=1}^n \frac{1}{2} \dot{\mathbf{X}}_i M_{ij} \dot{\mathbf{X}}_j \quad (4.15)$$

where M_{ij} is the n -vortex mass tensor given by

$$M_{ij} = \int d^2k \frac{q_i q_j}{J a^4 k^2 Q^4 r_v^2} (\hat{\mathbf{X}}_i \cdot \hat{\chi}_k) (\hat{\mathbf{X}}_j \cdot \hat{\chi}_k) e^{i\mathbf{k} \cdot (\mathbf{X}_i - \mathbf{X}_j)}$$

$$= \frac{\pi q_i q_j r_v^2}{J a^4} \begin{cases} \ln \frac{R_S}{r_{ij}} + \frac{1}{2} \left((\hat{\mathbf{X}}_i \cdot \hat{\mathbf{e}}_{ij})(\hat{\mathbf{X}}_j \cdot \hat{\mathbf{e}}_{ij}) \right. \\ \left. - (\hat{\mathbf{X}}_i \times \hat{\mathbf{e}}_{ij}) \cdot (\hat{\mathbf{X}}_j \times \hat{\mathbf{e}}_{ij}) \right), & i \neq j; \\ \ln \frac{R_S}{\sqrt{a^2 + r_v^2}}, & i = j. \end{cases} \quad (4.16)$$

where $\hat{\mathbf{e}}_{ij} = \frac{\mathbf{X}_i - \mathbf{X}_j}{|\mathbf{X}_i - \mathbf{X}_j|}$. This result is in agreement with that of Slonczewski⁵⁹. Slonczewski's calculation follows very closely that of section 3.4.

4.2.2 Radiation of magnons

In the previous section, we calculated the vortex inertial energy using second order perturbation theory. However, we only used the principle part of the integral. When evaluating the integral giving the second order perturbative energy shift¹⁵, to be careful in the divergent region $E_f \rightarrow E_i$, symbolically, we should write

$$\Delta E_i = \sum_f \frac{H_{if} H_{fi}}{E_i - E_f + i\epsilon} \quad (4.17)$$

and then take the limit $\epsilon \rightarrow 0$. But

$$\frac{1}{x + i\epsilon} = \frac{x}{x^2 + \epsilon^2} - \frac{i\epsilon}{x^2 + \epsilon^2}$$

The imaginary part approaches a δ -function as $\epsilon \rightarrow 0$ since

$$\int_{-\infty}^{\infty} dx \frac{\epsilon}{x^2 + \epsilon^2} = \pi \quad (4.18)$$

So then

$$\frac{1}{x + i\epsilon} = \text{principle value} \left(\frac{1}{x} \right) - i\pi\delta(x)$$

An imaginary part to the energy shift creates a decaying exponential factor in the time-dependent wave function

$$e^{-i(E/\hbar - i\gamma/2)t} = e^{iEt/\hbar} e^{-\gamma t/2}$$

and is hence interpreted as dissipation. The factor of 2 is there so that the probability $|\psi|^2$ decays as $e^{-\gamma t}$.

The rate of decay due to magnon emission is thus given symbolically by

$$\gamma = \sum_f \frac{2\pi}{\hbar} |H_{fi}|^2 \delta(E_f - E_i) \quad (4.19)$$

If we assume an initial state of no magnons the rate equation becomes

$$\gamma = 2\pi \int \frac{d^2ka^2}{(2\pi)^2} \frac{2\pi^2 Sq^2}{a^4 k Q^3 r_v^2} \left(\dot{\mathbf{X}} \cdot \hat{\chi}_k \right)^2 \delta \left(\frac{(\mathbf{P} - \hbar \mathbf{k})^2}{2M_v} + \hbar ckQ - \frac{P^2}{2M_v} \right) \quad (4.20)$$

Again, initially we have no inertial term, simplifying γ to

$$\gamma = 2\pi \int \frac{d^2ka^2}{(2\pi)^2} \frac{2\pi^2 Sq^2}{a^4 k Q^3 r_v^2} \left(\dot{\mathbf{X}} \cdot \hat{\chi}_k \right)^2 \delta(\hbar ckQ) \quad (4.21)$$

Rewriting the δ -function as

$$\delta(\hbar ckQ) = \frac{Q}{\hbar c(k^2 + Q^2)} \delta(k)$$

and the rate of dissipation becomes

$$\begin{aligned} \gamma &= \int \frac{d\chi_k}{2\pi} \frac{2\pi^2 q^2 r_v^2}{\hbar J a^4} \left(\dot{\mathbf{X}} \cdot \hat{\chi}_k \right)^2 \\ &= \frac{\pi^2 q^2 r_v^2 \dot{X}^2}{\hbar J a^4} \end{aligned} \quad (4.22)$$

Slonczewski⁵⁹ calculated microscopically a dissipation rate by extending the simple results of section 3.4 to include retardation effects of spin waves. However, to evaluate the far region perturbations, he assumes small frequencies which give a log divergent frequency dependent dissipation.

In our analysis, we used the same vortex-magnon coupling, although, accounting only for the $k = 0$ contribution. We find the full spectrum contributions in section 4.4.1 and will return to this comparison then.

Second order radiative corrections

Interestingly, the rate of emission using the finite mass of the vortex calculated in section 4.2.1 is actually considerably more important. This is reasonable since when we assume a finite mass, the vortex can lose kinetic energy by emitting a finite energy magnon.

Thus, with a finite M_v this time,

$$\gamma = 2\pi \int \frac{d^2k a^2}{(2\pi)^2} \frac{2\pi^2 S q^2}{a^4 k Q^3 r_v^2} \left(\dot{\mathbf{X}} \cdot \hat{\chi}_{\mathbf{k}} \right)^2 \delta \left(\frac{(\mathbf{P} - \hbar \mathbf{k})^2}{2M_v} + \hbar c k Q - \frac{P^2}{2M_v} \right) \quad (4.23)$$

Note that the vector potential component of the vortex momentum won't be included because in perturbation theory we assume that the vortex momentum is changed by changing speed, not position.

Let k_χ be a solution of the delta function condition as a function of emission angle, $\chi_{\mathbf{k}}$. The delta function can then be rewritten

$$\delta \left(\frac{\hbar^2}{2M_v} \left(k - \left(\frac{P}{\hbar} \cos \chi_{\mathbf{k}} - \frac{cM_v}{r_v \hbar} \right) \right)^2 - \frac{\hbar^2}{2M_v} \left(k_\chi - \left(\frac{P}{\hbar} \cos \chi_{\mathbf{k}} - \frac{cM_v}{r_v \hbar} \right) \right)^2 \right) \quad (4.24)$$

where we've approximated $Q \approx 1/r_v$, a reasonable approximation assuming small vortex velocities. Changing variables within the delta function to express it as $\delta(k - k_\chi)$, the integral becomes

$$\gamma = \frac{S\pi q^2 r_v}{a^2} \int d^2k \frac{\left(\dot{\mathbf{X}} \cdot \hat{\chi}_{\mathbf{k}} \right)^2}{k} \frac{M_v \delta(k - k_\chi)}{\hbar^2 \left| k_\chi - \left(\frac{P}{\hbar} \cos \chi_{\mathbf{k}} - \frac{cM_v}{r_v \hbar} \right) \right|} \quad (4.25)$$

Substituting for k_χ ,




Figure 4.3: The dissipation rate from perturbation theory; first assuming infinite mass and then adding corrections due to finite mass.

$$\begin{aligned}
\gamma &= \frac{S\pi q^2 r_v}{a^2} \int d^2k \frac{(\dot{\mathbf{X}} \cdot \hat{\chi}_{\mathbf{k}})^2}{k} \frac{M_v \delta(k - k_\chi)}{\hbar^2 \left| \frac{P}{\hbar} \cos \chi_{\mathbf{k}} - \frac{cM_v}{r_v \hbar} \right|} \\
&= \frac{S\pi q^2 r_v M_v \dot{X}^2}{\hbar P a^2} \int d\chi_{\mathbf{k}} \frac{\sin^2 \chi_{\mathbf{k}}}{\left| \cos \chi_{\mathbf{k}} - \frac{cM_v}{Pr_v} \right|} \\
&= \frac{2\pi q^2 S^2 J}{\hbar} \left(\frac{\pi}{2} + \sqrt{\tilde{P}^2 - 1} - \sin^{-1} \frac{1}{\tilde{P}} + \sqrt{\tilde{P}^2 - 1} \ln \left(\frac{\hbar \pi r_v}{R_S c M_v} \frac{\tilde{P}}{\tilde{P}^2 - 1} \right) \right)
\end{aligned} \tag{4.26}$$

where $\tilde{P} = \frac{Pr_v}{cM_v}$. To evaluate this last integral, an infrared cut-off had to be imposed: $k_{min} = \frac{2\pi}{R_S}$.

The discontinuity at $\tilde{P} = 1$ occurs when the vortex attains the minimum energy to overcome the “semi-gap” formed by the $Q = \sqrt{k^2 + 1/r_v^2}$ factor in the energy spectrum.

Note that this dissipation is in addition to that calculated in the previous section. We didn’t get both contributions here because we left out the $k = 0$ solution of the δ -function (4.24). See Figure 4.3 for a plot of these two contributions.

4.2.3 Zero point energy

Consider next the two magnon couplings, (4.3), arising from expanding the Hamiltonian about a stable vortex. Separate out the terms corresponding to the magnon Lagrangian expanded about a vacuum solution

$$\frac{S}{a^2} \dot{\vartheta} \vartheta - \frac{S^2 J}{2} \left((\nabla \vartheta)^2 + (\nabla \varphi)^2 + \frac{1}{r_v^2} \vartheta^2 \right)$$

and interpret those remaining as an interaction Hamiltonian

$$\begin{aligned} H_{int} = S \int \frac{d^2 r}{a^2} \left(-\dot{\varphi} (1 - \sin \theta_v) \vartheta + \frac{c}{2} \left(-\cos^2 \theta_v (\nabla \varphi)^2 + 2 \sin 2\theta_v \nabla \phi_v \cdot \nabla \varphi \vartheta \right. \right. \\ \left. \left. + \cos 2\theta_v (\nabla \phi_v)^2 \vartheta^2 - 2 \frac{\cos^2 \theta_v}{r_v^2} \vartheta^2 \right) \right) \end{aligned} \quad (4.27)$$

The only long range interaction term in H_{int} (i.e. that doesn't decay exponentially) is the $\sin^2 \theta_v$ portion of the fourth term. Fourier transforming the ϑ factors according to (2.5), this long range term becomes

$$\begin{aligned} H_{int} &= -\frac{Sca^2}{2} \int d^2 r (\nabla \phi_v)^2 \int \frac{d^2 k}{(2\pi)^2} e^{-i\mathbf{k} \cdot \mathbf{r}} e^{-i\mathbf{k} \cdot \mathbf{X}} \vartheta_{\mathbf{k}} \int \frac{d^2 k'}{(2\pi)^2} e^{i\mathbf{k}' \cdot \mathbf{r}} e^{i\mathbf{k}' \cdot \mathbf{X}} \vartheta_{-\mathbf{k}'} \\ &= -Sca^2 q^2 \pi \int dr \frac{J_0(|\mathbf{k} - \mathbf{k}'|r)}{r} \int \frac{d^2 k}{(2\pi)^2} \int \frac{d^2 k'}{(2\pi)^2} e^{-i(\mathbf{k} - \mathbf{k}') \cdot \mathbf{X}} \vartheta_{\mathbf{k}} \vartheta_{-\mathbf{k}'} \end{aligned} \quad (4.28)$$

This integral over r diverges in the short range. However, the original term actually changes sign as $r \rightarrow 0$ so that the analysis is invalid into the core region anyway and must be cut-off. We define a form factor $\mathcal{F}(\kappa) = e^{-i\kappa \cdot \mathbf{X}} \int \frac{J_0(\kappa r)}{r} dr$ where $\kappa = \mathbf{k} - \mathbf{k}'$.

Expressing this term in the language of quantized magnons (see section 2.1) gives a Fourier transformed version

$$\begin{aligned}
H_{int}^{(1)} &= -S^2 J q^2 \pi \int \frac{a^2 d^2 k}{(2\pi)^2} \frac{a^2 d^2 \kappa}{(2\pi)^2} \mathcal{F}(\kappa) \vartheta_{\mathbf{k}+\kappa} \vartheta_{-\mathbf{k}} \\
&= \frac{\hbar S J q^2 a^4 \pi}{2} \int \frac{d^2 k}{(2\pi)^2} \frac{d^2 \kappa}{(2\pi)^2} \mathcal{F}(\kappa) \left(\frac{k \kappa}{Q \sqrt{\kappa^2 + 1/r_v^2}} \right)^{\frac{1}{2}} \\
&\quad \times \left(a_{\mathbf{k}+\kappa}^\dagger - a_{-(\mathbf{k}+\kappa)} \right) \left(a_{-\mathbf{k}}^\dagger - a_{\mathbf{k}} \right) \\
&= \frac{\hbar S J q^2 a^4 \pi}{2} \int \frac{d^2 k}{(2\pi)^2} \frac{d^2 \kappa}{(2\pi)^2} \mathcal{F}(\kappa) \left(\frac{k \kappa}{Q \sqrt{\kappa^2 + 1/r_v^2}} \right)^{\frac{1}{2}} \\
&\quad \times \left(a_{\mathbf{k}+\kappa}^\dagger a_{-\mathbf{k}}^\dagger + a_{-(\mathbf{k}+\kappa)} a_{\mathbf{k}} - a_{\mathbf{k}+\kappa}^\dagger a_{\mathbf{k}} - a_{\mathbf{k}}^\dagger a_{\mathbf{k}+\kappa} - (2\pi)^2 \frac{\delta^2(\kappa)}{a^2} \right)
\end{aligned} \tag{4.29}$$

The a and $a^\dagger$ terms above correspond, respectively, to the case of two magnons being created in opposite directions, two magnons incoming from opposite directions being annihilated, a magnon given a momentum boost of κ , and the last combination removes momentum κ from an existing magnon. The last term gives the zero point energy shift

$$\begin{aligned}
\Delta E &= -\frac{\hbar c q^2 \pi}{2} \ln \frac{R_s}{r_v} \int \frac{d^2 k}{(2\pi)^2} \frac{k}{Q} \\
&= -\frac{\hbar c q^2}{4} \ln \frac{R_s}{r_v} \int dk \frac{k^2}{Q} \\
&= -\frac{\hbar c q^2}{4} \ln \frac{R_s}{r_v} \left(\frac{\sqrt{r_v^2 + a^2}}{2a^2 r_v} - \frac{1}{2r_v^2} \ln \left(\frac{r_v + \sqrt{r_v^2 + a^2}}{a} \right) \right)
\end{aligned} \tag{4.30}$$

as found before in section 3.5.

4.3 Vortex influence functional

In this section, we develop the effective dynamics of the magnetic vortex using path integration. The temperature is introduced by assuming as an initial condition that the magnons are in thermal equilibrium. They are of course allowed to evolve out of equilibrium when interactions with the vortex are considered.

Populating the magnons at a temperature τ , we have a density matrix describing them given by equation (2.27). To describe the effective dynamics of the reduced density matrix for the vortex, we trace out the magnon degrees of freedom from the full vortex-magnon density matrix using the Feynman-Vernon influence functional formalism¹⁷. Before proceeding, consider the simple case of

a central coordinate $x(t)$ coupled to a bath of simple harmonic oscillators $r_i(t)$ with frequencies ω_i . This introduces the influence functional formalism and the interpretation of results for our own magnetic system.

Separate the Lagrangian describing a coordinate $x(t)$ coupled linearly to a set of harmonic oscillators r_i as

$$\mathcal{L} = \mathcal{L}_x[x(t)] + \mathcal{L}_r[r_i] + \mathcal{L}_{int}[x(t), r_i(t)] \quad (4.31)$$

where $\mathcal{L}_x[x(t)]$ describes subsystem $x(t)$, $\mathcal{L}_r[r_i]$ describes the environmental modes and $\mathcal{L}_{int}[x(t), r_i(t)]$ describes the couplings between the two systems.

Assume a general Lagrangian $\mathcal{L}_x[x(t)]$ for the central coordinate, a simple harmonic Lagrangian in r_i

$$\mathcal{L}_r[r_i] = \sum_i \frac{1}{2} \dot{r}_i^2 + \frac{\omega_i^2}{2} r_i^2 \quad (4.32)$$

and for the interacting Lagrangian, assume linear couplings

$$\mathcal{L}_{int}[x(t), r_i(t)] = \sum_i C_i x(t) r_i(t) \quad (4.33)$$

Generally, the dynamics of the two subsystems become entangled which is conveniently described within the density matrix formalism.

The density matrix of the complete system in operator form evolves from initial state $\rho(0)$ according to

$$\rho(T) = \exp - \frac{iHT}{\hbar} \rho(0) \exp \frac{iHT}{\hbar} \quad (4.34)$$

Alternatively, in the coordinate representation,

$$\begin{aligned} \rho(x, r_i, T; y, q_i, 0) &= \langle x, r_i | \rho(T) | y, q_i \rangle \\ &= \int dx' dy' dr'_i dq'_i \langle x, r_i | \exp - \frac{iHT}{\hbar} | x', r'_i \rangle \\ &\quad \times \langle x', r'_i | \rho(0) | y', q'_i \rangle \langle y', q'_i | \exp \frac{iHT}{\hbar} | y, q_i \rangle \end{aligned} \quad (4.35)$$

Expanding each propagator as a path integral, noting

$$\begin{aligned} \langle x, r_i | \exp - \frac{iHT}{\hbar} | x', r'_i \rangle &= \int_{x'}^x \mathcal{D}[x(t)] \int_{r'_i}^{r_i} \mathcal{D}[r_i(t)] \exp \frac{i}{\hbar} \mathcal{S}[x(t), r_i(t)] \\ \langle x, r_i | \exp - \frac{iHT}{\hbar} | x', r'_i \rangle &= \int_{y'}^y \mathcal{D}[y(t)] \int_{q'_i}^{q_i} \mathcal{D}[q_i(t)] \exp - \frac{i}{\hbar} \mathcal{S}[y(t), q_i(t)] \end{aligned}$$

the density matrix at time T becomes

$$\begin{aligned} \rho(x, r_i; y, q_i; T) &= \int dx' dy' dr'_i dq'_i \int_{x'}^x \mathcal{D}[x(t)] \int_{r'_i}^{r_i} \mathcal{D}[r_i(t)] \exp \frac{i}{\hbar} \mathcal{S}[x(t), r_i(t)] \\ &\quad \times \langle x', r'_i | \rho(0) | y', q'_i \rangle \int_{y'}^y \mathcal{D}[y(t)] \int_{q'_i}^{q_i} \mathcal{D}[q_i(t)] \exp -\frac{i}{\hbar} \mathcal{S}[y(t), q_i(t)] \end{aligned}$$

However, suppose we're only interested in the dynamics of the subsystem $x(t)$, regardless of the specific behaviour of the harmonic oscillator subsystems. To eliminate these variables, perform the trace over the $\{r_i\}$ variables to obtain the so-called reduced density operator

$$\begin{aligned} \tilde{\rho}(x; y; T) &= \int dr_i \int dx' dy' dr'_i dq'_i \int_{x'}^x \mathcal{D}[x(t)] \int_{r'_i}^{r_i} \mathcal{D}[r_i(t)] \exp \frac{i}{\hbar} \mathcal{S}[x(t), r_i(t)] \\ &\quad \times \langle x', r'_i | \rho(0) | y', q'_i \rangle \int_{y'}^y \mathcal{D}[y(t)] \int_{q'_i}^{q_i} \mathcal{D}[q_i(t)] \exp -\frac{i}{\hbar} \mathcal{S}[y(t), q_i(t)] \end{aligned}$$

Assume that the $t = 0$ density matrix is separable in the two subsystems, i.e. that they are initially disentangled and

$$\rho(x, r_i; y, q_i; 0) = \rho_x(x, y; 0) \rho_r(r_i, q_i; 0) \quad (4.36)$$

Further, assume that the simple harmonic oscillators are initially in thermal equilibrium so that $\rho_r(r_i, q_i; t = 0)$ is given by¹⁶

$$\rho_r(r_i, q_i; 0) = \prod_i \sqrt{\frac{m\omega_i}{2\pi\hbar \sinh \hbar\omega_i\beta}} \exp -\frac{m\omega}{2\hbar \sinh \hbar\omega_i\beta} ((r_i^2 + q_i^2) \cosh \hbar\omega_i\beta - 2r_i q_i)$$

The reduced density matrix is then expressible as

$$\tilde{\rho}(x; y; t = T) = \int dx' \int dy' J(x, y, T; x', y', 0) \rho_x(x', y', 0) \quad (4.37)$$

where

$$J(x, y, T; x', y', 0) = \int_{x'}^x \mathcal{D}[x(t)] \int_{y'}^y \mathcal{D}[y(t)] \exp \frac{i}{\hbar} (\mathcal{S}_x[x(t)] - \mathcal{S}_x[y(t)]) \mathcal{F}[x(t), y(t)] \quad (4.38)$$

is the propagator for the density operator and

$$\begin{aligned}
F[x(t), y(t)] &= \int dr_i dr'_i dq'_i \rho_r(r'_i, q'_i, 0) \int_{r'_i}^{r_i} \mathcal{D}[r_i(t)] \int_{q'_i}^{q_i} \mathcal{D}[q_i(t)] \\
&\times \exp \frac{i}{\hbar} (S_r[r_i(t)] + S_{int}[r_i(t), x(t)] - S_r[q_i(t)] - S_{int}[q_i(t), y(t)])
\end{aligned} \tag{4.39}$$

is the influence functional¹⁷. Evaluating this for the central coordinate $x(t)$ coupled linearly to a set of environmental modes described by simple harmonic oscillators with spectrum $\omega_i(t)$

$$F[x, y] = \exp -\frac{1}{\hbar} \int_0^T dt \int_0^t ds (x(t) - y(t)) (\alpha(t-s)x(s) - \alpha^*(t-s)y(s)) \tag{4.40}$$

where

$$\alpha(t-s) = \sum_i \frac{C_i^2}{2m\omega_i} \left(\exp -i\omega_i(t-s) + \frac{2 \cos \omega_i(t-s)}{\exp \hbar\omega_i\beta - 1} \right) \tag{4.41}$$

where C_i are the linear coupling parameters.

4.3.1 Quantum Brownian motion

Caldeira and Leggett⁵ interpret the influence functional result as quantized damped dynamics. The problem of quantizing Brownian motion was not entirely understood. Their idea of coupling to a bath of oscillators to achieve Brownian motion (which, of course, from there is easily quantizable) was one of many proposed in the 1980's and 90's. The classical equation of motion for Brownian motion, the Langevin equation, is

$$m\ddot{x} + \eta\dot{x} + V'(x) = F(t) \tag{4.42}$$

where m is the mass of the particle, η is a damping constant, $V(x)$ is the potential acting on the particle and $F(t)$ is the fluctuating force. This force obeys

$$\begin{aligned}
\langle F(t) \rangle &= 0 \\
\langle F(t)F(t') \rangle &= 2\eta kT \delta(t-t')
\end{aligned} \tag{4.43}$$

where $\langle \rangle$ denote statistical averaging.

With such a force, the propagator of the density matrix of system x is given by

$$J(x, y, t; x', y', 0) = \int \mathcal{D}[x] \mathcal{D}[y] \exp \frac{i}{\hbar} \left(\mathcal{S}[x] - \mathcal{S}[y] + \int_0^t d\tau (x(\tau) - y(\tau)) F(\tau) \right)$$

Assuming that the fluctuating force $F(t)$ has the probability distribution functional $P[F(\tau)]$ of different histories $F(\tau)$, the averaged density matrix propagator becomes

$$J(x, y, t; x', y', 0) = \int \mathcal{D}[x] \mathcal{D}[y] \mathcal{D}[F] P[F(\tau)] \exp \frac{i}{\hbar} (\mathcal{S}[x] - \mathcal{S}[y] + \int_0^t d\tau (x(\tau) - y(\tau)) F(\tau)) \quad (4.44)$$

We can perform the path integration over $F(\tau)$ if we assume $P[F(\tau)]$ is a Gaussian distribution, yielding

$$J(x, y, t; x', y', 0) = \int \mathcal{D}[x] \mathcal{D}[y] \exp \frac{i}{\hbar} (\mathcal{S}[x] - \mathcal{S}[y]) \times \exp -\frac{1}{\hbar^2} \int_0^t \int_0^\tau d\tau ds (x(\tau) - y(\tau)) A(\tau - s) (x(s) - y(s)) \quad (4.45)$$

where $A(\tau - s)$ is the correlation of forces, $\langle F(\tau) F(s) \rangle$.

The real exponentiated term in the influence functional is

$$\exp -\frac{1}{\hbar} \int_0^t \int_0^\tau d\tau ds (x(\tau) - y(\tau)) \alpha_R(\tau - s) (x(s) - y(s)) \quad (4.46)$$

where

$$\alpha_R(\tau - s) = \sum_i \frac{C_i^2}{2m\omega_i} \coth \frac{\hbar\omega_i}{2k_B T} \cos \omega_i(\tau - s) \quad (4.47)$$

where C_i denotes the coupling coefficient to the i th environmental mode. Assuming instead a continuum of k states with density

$$\rho_D(\omega) C^2(\omega) = \begin{cases} \frac{2m\eta\omega^2}{\pi}, & \omega < \Omega; \\ 0, & \omega > \Omega. \end{cases} \quad (4.48)$$

the influence functional result becomes in a high temperature limit ($\coth \frac{\hbar\omega}{2kT} \rightarrow \frac{2kT}{\hbar\omega}$)

$$\hbar\alpha_R(\tau - s) = \langle F(\tau) F(s) \rangle = 2\eta kT \frac{\sin \Omega(\tau - s)}{\pi(\tau - s)} \quad (4.49)$$

which tends to (4.43) in the limit $\Omega \rightarrow \infty$.

The imaginary phase term in the influence functional is manipulated to give an x^2 frequency shift which renormalizes the external potential. In addition to this, there is a new action term corresponding to a damping force

$$\Delta\mathcal{S} = - \int_0^t dt M \gamma (x\dot{x} - y\dot{y} + x\dot{y} - y\dot{x}) \quad (4.50)$$

Note that the forward and backward paths are interacting so that the new effective action is coupled in $x(t)$ and $y(t)$.

The relaxation constant γ is

$$\gamma = \frac{\eta}{2M} \quad (4.51)$$

where the damping constant η is dependent on the density of states of the environmental modes. For our treatment where the environmental modes are magnons, we know explicitly the magnon density of states, going as $\frac{Q}{k^2+Q^2}$ (recall $\omega(k) = ckQ$) rather than ω^2 as assumed above, so that our analysis does not simplify to a frequency independent damping function.

Castro Neto and Caldeira⁴² consider the problem of a central coordinate coupled linearly to a set of oscillators; however, as opposed to the Caldeira and Leggett⁵ problem, the central system, $X(t)$, is a solution in the same medium as the set of oscillators. Hence, as in our problem, there is no linear coupling with position, but instead, we find a linear coupling $C_i r_i \dot{X}(t)$ between oscillators $\{r_i\}$ and the velocity.

They simplify their results by assuming a Born approximation. Although they lose the resulting frequency dependent motion, they do find that the damping coefficients and correlation integrals now possess memory effects. We will discuss these issues after results have been simplified for our vortex-magnon system.

4.3.2 Semiclassical solution of perturbed magnons

Before evaluating the influence functional, we first need the propagator of the magnon system perturbed by the vortex presence. The effect of important perturbing terms have been discussed already using perturbation theory. The one magnon coupling endows the vortex with an effective mass and makes the vortex motion dissipative by radiating magnons. The leading two magnon coupling provides an overall zero-point energy shift to the vortex-magnon system that is associated to the quantized vortex. Although the two magnon couplings, or indeed any of the many magnon couplings, may give more significant dissipation, we neglect these contributions in this treatment. In the influence functional, the forward and backward paths have cancelling zero point energy shifts and hence we will ignore entirely the many magnon couplings.

Treat the disturbance of a magnetic vortex centered at $\mathbf{X}(t)$ with vorticity q and polarization p by the magnons via the one magnon coupling, (4.6),

$$\mathcal{L}_{int} = -S \int \frac{d^2 r}{a^2} \dot{\mathbf{X}} \cdot \nabla \phi_v \sin \theta_v \vartheta \quad (4.52)$$

We must evaluate the propagator for the system of magnons, again in the φ basis

$$\langle \varphi | \exp -\frac{i\mathcal{H}t}{\hbar} | \varphi' \rangle = \int_{\varphi'}^{\varphi} \mathcal{D}[\varphi] \mathcal{D}[\vartheta] \exp \frac{i}{\hbar} \int_0^T dt (\mathcal{L}_m + \mathcal{L}_{int}) \quad (4.53)$$

where $\mathbf{X}(t)$ is considered now an externally controlled parameter.

Introduce the plane wave decomposition (2.5) so that the action becomes

$$\begin{aligned} \mathcal{S}_{m+int}[\varphi, \vartheta] = S \int \frac{a^2 d^2 k}{(2\pi)^2} \int_0^T dt & \left(\dot{\varphi}_{\mathbf{k}} \vartheta_{-\mathbf{k}} - \frac{c}{2} (k^2 \varphi_{\mathbf{k}} \varphi_{-\mathbf{k}} + Q^2 \vartheta_{\mathbf{k}} \vartheta_{-\mathbf{k}}) \right. \\ & \left. - \int \frac{d^2 r}{a^2} e^{-i\mathbf{k} \cdot \mathbf{r}} \dot{\mathbf{X}} \cdot \nabla \phi_v \sin \theta_v \vartheta_{\mathbf{k}} \right) \end{aligned} \quad (4.54)$$

The equations of motion are modified by a force term, that, for simplicity, we denote as

$$\begin{aligned} f_{\mathbf{k}}[\mathbf{X}] &= \int \frac{d^2 r}{a^2} e^{i\mathbf{k} \cdot \mathbf{r}} \dot{\mathbf{X}} \cdot \nabla \phi_v \sin \theta_v \\ &= \frac{2i\pi q}{a^2} \frac{\dot{\mathbf{X}} \cdot \hat{\varphi}_{\mathbf{k}}}{kQr_v} e^{i\mathbf{k} \cdot \mathbf{X}} \end{aligned} \quad (4.55)$$

and become

$$\begin{pmatrix} ck^2 & \frac{\partial}{\partial t} \\ -\frac{\partial}{\partial t} & cQ^2 \end{pmatrix} \begin{pmatrix} \varphi_{\mathbf{k}}^{cl} \\ \vartheta_{\mathbf{k}}^{cl} \end{pmatrix} = \begin{pmatrix} 0 \\ -f_{\mathbf{k}}[\mathbf{X}] \end{pmatrix} \quad (4.56)$$

The solution with boundary conditions $\varphi_{\mathbf{k}}(0) = \varphi'_{\mathbf{k}}$ and $\varphi_{\mathbf{k}}(T) = \varphi_{\mathbf{k}}$ is

$$\begin{aligned} \begin{pmatrix} \varphi_{\mathbf{k}}^{cl} \\ \vartheta_{\mathbf{k}}^{cl} \end{pmatrix} &= \frac{\varphi_{\mathbf{k}} - \int_t^T ds \cos \omega_k (T-s) f_{\mathbf{k}}[\mathbf{X}]}{\sin \omega_k T} \begin{pmatrix} \sin \omega_k t \\ \frac{k}{Q} \cos \omega_k t \end{pmatrix} \\ &+ \frac{\varphi'_{\mathbf{k}} + \int_0^t ds \cos \omega_k s f_{\mathbf{k}}[\mathbf{X}]}{\sin \omega_k T} \begin{pmatrix} \sin \omega_k (T-t) \\ -\frac{k}{Q} \cos \omega_k (T-t) \end{pmatrix} \end{aligned} \quad (4.57)$$

Substituting this solution into the action gives the classical contribution

$$\begin{aligned} \mathcal{S}_{m+int}^{cl}(k) = & \frac{Sk}{2Q \sin \omega_k T} \left((\varphi'_k \varphi'_{-k} + \varphi_k \varphi_{-k}) \cos \omega_k T - 2\varphi_k \varphi'_{-k} \right. \\ & - 2\varphi_k \int_0^T dt \cos \omega_k t f_{-k}[\mathbf{X}(t)] + 2\varphi'_k \int_0^T dt \cos \omega_k (T-t) f_{-k}[\mathbf{X}(t)] \\ & \left. + 2 \int_0^T dt \int_0^t ds \cos \omega_k (T-t) \cos \omega_k s f_k[\mathbf{X}(t)] f_{-k}[\mathbf{X}(s)] \right) \end{aligned} \quad (4.58)$$

The quantum fluctuations introduce a pre-factor given by solving the relevant Jacobi equation

$$\begin{pmatrix} ck^2 & \frac{\partial}{\partial t} \\ -\frac{\partial}{\partial t} & cQ^2 \end{pmatrix} \begin{pmatrix} \varphi(t) \\ \vartheta(t) \end{pmatrix} = 0 \quad (4.59)$$

with initial conditions $\varphi_k(0) = 0$ and $S\vartheta(0) = 1$. The determinant is given as $i\varphi(T)$. Combining the pre-factor $(\det)^{-1}$ with path integration measure factors, give the overall result, (2.18),

$$\sqrt{\frac{Sk}{2\pi i \hbar Q \sin \omega_k T}} \quad (4.60)$$

The final propagator is

$$\langle \varphi | \exp -\frac{i(H_{m+int})t}{\hbar} | \varphi' \rangle = \prod_{\mathbf{k}} \sqrt{\frac{Sk}{2\pi i \hbar Q \sin \omega_k T}} \exp \frac{i}{\hbar} \int \frac{a^2 d^2 k}{(2\pi)^2} \mathcal{S}_{m+int}^{cl}(k) \quad (4.61)$$

4.3.3 Evaluating the influence functional

Substituting the semiclassical solutions to the two path integrals and for the thermal equilibrium density matrix, the problem is reduced to three regular gaussian integrals, ignoring pre-factors, which cancel anyway after all integrals when the density matrix is properly normalized,

$$\begin{aligned}
F[\mathbf{X}, \mathbf{Y}] = & \prod_{\mathbf{k}} \int d\varphi_{\mathbf{k}} d\varphi'_{\mathbf{k}} d\tilde{\varphi}'_{\mathbf{k}} \\
& \exp \left[-\frac{Sk}{2\hbar Q \sinh \hbar\omega_k \beta} \left((\varphi'_{\mathbf{k}} \varphi'_{-\mathbf{k}} + \tilde{\varphi}'_{\mathbf{k}} \tilde{\varphi}'_{-\mathbf{k}}) \cosh \hbar\omega_k \beta - 2\tilde{\varphi}'_{\mathbf{k}} \varphi'_{-\mathbf{k}} \right) \right] \\
& \exp \left[\frac{iSk}{2\hbar Q \sin \omega_k T} \left((\varphi'_{\mathbf{k}} \varphi'_{-\mathbf{k}} + \varphi_{\mathbf{k}} \varphi_{-\mathbf{k}}) \cos \omega_k T - 2\varphi_{\mathbf{k}} \varphi'_{-\mathbf{k}} \right. \right. \\
& \quad - 2\varphi_{\mathbf{k}} \int_0^T dt \cos \omega_k t f_{-\mathbf{k}}[\mathbf{X}(t)] + 2\varphi'_{\mathbf{k}} \int_0^T dt \cos \omega_k (T-t) f_{-\mathbf{k}}[\mathbf{X}(t)] \\
& \quad \left. \left. + 2 \int_0^T dt \int_0^t ds \cos \omega_k (T-t) \cos \omega_k s f_{\mathbf{k}}[\mathbf{X}(t)] f_{-\mathbf{k}}[\mathbf{X}(s)] \right) \right] \\
& \exp \left[-\frac{iSk}{2\hbar Q \sin \omega_k T} \left((\tilde{\varphi}'_{\mathbf{k}} \tilde{\varphi}'_{-\mathbf{k}} + \varphi_{\mathbf{k}} \varphi_{-\mathbf{k}}) \cos \omega_k T - 2\tilde{\varphi}'_{\mathbf{k}} \varphi_{-\mathbf{k}} \right. \right. \\
& \quad - 2\varphi_{\mathbf{k}} \int_0^T dt \cos \omega_k t f_{-\mathbf{k}}[\mathbf{Y}(t)] + 2\tilde{\varphi}'_{\mathbf{k}} \int_0^T dt \cos \omega_k (T-t) f_{-\mathbf{k}}[\mathbf{Y}(t)] \\
& \quad \left. \left. + 2 \int_0^T dt \int_0^t ds \cos \omega_k (T-t) \cos \omega_k s f_{\mathbf{k}}[\mathbf{Y}(t)] f_{-\mathbf{k}}[\mathbf{Y}(s)] \right) \right]
\end{aligned} \tag{4.62}$$

where the $\prod_{\mathbf{k}}$ applies to everything (and hence implies integrals over $\mathbf{k}$ within exponentials).

Performing these integrals mimics very closely the calculations for the analogous problem of a central coordinate $x(t)$ coupled linearly to the position coordinate of a system of simple harmonic oscillators¹⁷. In fact, the final expression is the same, with the same substitution $m\omega \rightarrow Sk/Q$ found earlier in evaluating the magnon propagator (see section 2.2),

$$\begin{aligned}
F[\mathbf{X}, \mathbf{Y}] = & \exp -\frac{1}{\hbar} \int \frac{a^2 d^2 k}{(2\pi)^2} \int_0^T dt \int_0^t ds (f_{\mathbf{k}}[\mathbf{X}(t)] - f_{\mathbf{k}}[\mathbf{Y}(t)]) \\
& (\alpha_k(t-s) f_{-\mathbf{k}}[\mathbf{X}(s)] - \alpha_k^*(t-s) f_{-\mathbf{k}}[\mathbf{Y}(s)])
\end{aligned} \tag{4.63}$$

where

$$\alpha_k(t-s) = \frac{Sk}{2Q} \left(e^{-i\omega_k(t-s)} + \frac{2 \cos \omega_k(t-s)}{e^{\hbar\omega_k \beta} - 1} \right) \tag{4.64}$$

The propagator of the density operator can be written as

$$\begin{aligned}
J(\mathbf{X}, \mathbf{Y}, T; \mathbf{X}', \mathbf{Y}', 0) = & \int_{\mathbf{X}}^{\mathbf{X}'} \mathcal{D}\mathbf{X} \int_{\mathbf{Y}}^{\mathbf{Y}'} \mathcal{D}\mathbf{Y} \exp \frac{i}{\hbar} \left[\mathcal{S}_v[\mathbf{X}] - \mathcal{S}_v[\mathbf{Y}] - \right. \\
& \int \frac{a^2 d^2 k}{(2\pi)^2} \int_0^T dt \int_0^t ds \left((f_{\mathbf{k}}[\mathbf{X}(t)] - f_{\mathbf{k}}[\mathbf{Y}(t)]) \alpha_k^I(t-s) (f_{-\mathbf{k}}[\mathbf{X}(s)] + f_{-\mathbf{k}}[\mathbf{Y}(s)]) \right. \\
& \left. \left. - i (f_{\mathbf{k}}[\mathbf{X}(t)] - f_{\mathbf{k}}[\mathbf{Y}(t)]) \alpha_k^R(t-s) (f_{-\mathbf{k}}[\mathbf{X}(s)] - f_{-\mathbf{k}}[\mathbf{Y}(s)]) \right) \right]
\end{aligned} \tag{4.65}$$

where $\alpha_k^R(t-s)$ and $\alpha_k^I(t-s)$ are the real and imaginary parts of $\alpha_k(t-s)$

$$\begin{aligned}
\alpha_k^R(t-s) &= \frac{Sk}{2Q} \cos \omega_k(t-s) \coth \frac{\hbar \omega_k \beta}{2} \\
\alpha_k^I(t-s) &= -\frac{Sk}{2Q} \sin \omega_k(t-s)
\end{aligned} \tag{4.66}$$

Ordinarily, we would extract a spectral function $\mathcal{J}(\omega, T)$ from this result that gives the frequency waiting of the functions α . For example, the results of Caldeira and Leggett can be re-expressed as

$$\begin{aligned}
\alpha_k^R(t-s) &= \int \frac{d\omega}{\pi} \mathcal{J}(\omega, T) \cos \omega(t-s) \coth \frac{\hbar \omega \beta}{2} \\
\alpha_k^I(t-s) &= -\int \frac{d\omega}{\pi} \mathcal{J}(\omega, T) \sin \omega(t-s)
\end{aligned} \tag{4.67}$$

In our case, however, we must first integrate over the angular dependence of $\mathbf{k}$. This, however, gives the sum of two terms with Bessel function factors of order 0 and 2, themselves dependent on the wavenumber k and the coordinate path $\mathbf{X}(t)$. In order to define a spectral function, we would have to disentangle the $t-s$ and k behaviours, which, with the additional $J_i(k|\mathbf{X}(t) - \mathbf{X}(s)|)$ factors is rather involved.

4.3.4 Interpreting the imaginary part

The one magnon coupling treated perturbatively endows the vortex with an effective mass and introduces dissipation. In the influence formalism, we expect to obtain terms in the effective action of the forward/backward paths interpretable as particle-like inertial terms. Dissipation arises due to fluctuating forces inflicted by scattered magnons on the vortex. We expect the fluctuating forces to be accompanied by corresponding damping forces.

Our one magnon term couples to the vortex velocity and not position as treated by Caldeira and Leggett. This is because the vortex is a solution itself of the

system, so that all first order variations vanish. The velocity term survives because the vortex is to zeroth order a stationary solution. The potential renormalization found earlier going like x^2 should here appear as a shift $\sim \dot{X}(t)^2$, or an inertial term from which we can deduce an effective vortex mass.

Substituting for $f_{\mathbf{k}}$ into the imaginary term yields the phase, including the additional minus sign in (4.65),

$$\Phi = -\frac{Sq^2}{2a^2} \int d^2k \int_0^T dt \int_0^t ds \left(\dot{\mathbf{X}}(t) e^{i\mathbf{k} \cdot \mathbf{X}(t)} - \dot{\mathbf{Y}}(t) e^{i\mathbf{k} \cdot \mathbf{Y}(t)} \right) \cdot \hat{\varphi}_{\mathbf{k}} \frac{\sin \omega_k(t-s)}{kQ^3 r_v^2} \\ \left(\dot{\mathbf{X}}(s) e^{-i\mathbf{k} \cdot \mathbf{X}(s)} + \dot{\mathbf{Y}}(s) e^{-i\mathbf{k} \cdot \mathbf{Y}(s)} \right) \cdot \hat{\varphi}_{\mathbf{k}} \quad (4.68)$$

where we define the phase angles via

$$F = \exp \left(\frac{-i}{\hbar} (\Phi - i\Gamma) \right) \quad (4.69)$$

Performing first the integral over χ_k from 0 to π (refer to an identical calculation in the perturbation calculation of multi-vortex mass corrections in section 4.2.1) to leading order yields, for the X^2 term only for conciseness (the other factors have the same form),

$$\Phi = -\frac{\pi Sq^2}{2a^2} \int dk \int_0^T dt \int_0^t ds \frac{\sin \omega_k(t-s)}{Q^3 r_v^2} \left(\dot{\mathbf{X}}(t) \cdot \dot{\mathbf{X}}(s) J_0(k|\mathbf{X}(t) - \mathbf{X}(s)|) \right. \\ \left. + \dot{X}(t) \dot{X}(s) \left((\hat{\mathbf{X}}(t) \cdot \hat{\mathbf{e}}_{\Delta})(\hat{\mathbf{X}}(s) \cdot \hat{\mathbf{e}}_{\Delta}) \right. \right. \\ \left. \left. - (\hat{\mathbf{X}}(t) \times \hat{\mathbf{e}}_{\Delta}) \cdot (\hat{\mathbf{X}}(s) \times \hat{\mathbf{e}}_{\Delta}) \right) J_2(k|\mathbf{X}(t) - \mathbf{X}(s)|) \right) + \text{etc.} \quad (4.70)$$

where $\hat{\mathbf{e}}_{\Delta}$ denotes the unit vector connecting $\mathbf{X}(t)$ and $\mathbf{X}(s)$.

Integrate by parts in $t-s$ to get two terms, one with two time derivatives in $X(s)$ and another with a single time derivative in $X(s)$. Note, we ignore the derivatives of the Bessel functions since the extra factor of k makes these higher order corrections and we assume the vortex curves slowly to ignore derivatives of the unit vectors. The boundary terms from the integration by parts are zero for $t=s$ and otherwise unimportant (they don't contribute to the equations of motion, being just boundary dependant). Finally, we have

$$\Phi = -\frac{Sq^2\pi}{2a^2} \int dk k dt ds \left(\frac{\sin \omega_0(t-s)}{kQ^3 r_v^2} \ddot{\mathbf{X}}(s) - \frac{c \cos \omega_0(t-s)}{Q^2 r_v^2} \dot{\mathbf{X}}(s) \right) \cdot \mathbf{X}(t) \\ \times J_0(k|\mathbf{X}(t) - \mathbf{X}(s)|) + \left(\frac{\sin \omega_0(t-s)}{kQ^3 r_v^2} \ddot{X}(s) - \frac{c \cos \omega_0(t-s)}{Q^2 r_v^2} \dot{X}(s) \right) \\ \times \left[\hat{\mathbf{X}}(s) \cdot \left(\hat{\mathbf{e}}_{\Delta}(\mathbf{X}(t) \cdot \hat{\mathbf{e}}_{\Delta}) - \hat{\mathbf{e}}_{\perp}(\mathbf{X}(t) \cdot \hat{\mathbf{e}}_{\perp}) \right) \right] J_2(k|\mathbf{X}(t) - \mathbf{X}(s)|) + \text{etc.} \quad (4.71)$$

where $\hat{\mathbf{e}}_\perp$ is a unit vector perpendicular to $\hat{\mathbf{e}}_\Delta$.

Split this integral into the sine and cosine portions, $\Phi = \Phi_s + \Phi_c$. Consider first the sine integrals. Integrate by parts again in $t - s$. The non-zero boundary terms are

$$\begin{aligned}\Phi_s^{BC} &= \frac{Sq^2\pi}{2a^2} \int dk \int_0^T dt \frac{1}{ckQ^4r_v^2} \left(\ddot{\mathbf{X}}(t) \cdot \mathbf{X}(t) - \ddot{\mathbf{Y}}(t) \cdot \mathbf{Y}(t) \right) \\ &= \frac{Sq^2\pi r_v^2}{2a^2c} \int_0^T dt \left(\ddot{\mathbf{X}}(t) \cdot \mathbf{X}(t) - \ddot{\mathbf{Y}}(t) \cdot \mathbf{Y}(t) \right) \ln \frac{R_S}{\sqrt{a^2 + r_v^2}}\end{aligned}\quad (4.72)$$

or equivalently,

$$\Phi_s^{BC} = -\frac{Sq^2\pi r_v^2}{2a^2c} \int_0^T dt \left(\dot{\mathbf{X}}^2(t) - \dot{\mathbf{Y}}^2(t) \right) \ln \frac{R_S}{\sqrt{a^2 + r_v^2}} \quad (4.73)$$

where we've again split the sine integral into $\Phi_s = \Phi_s^{BC} + \Phi_s^{int}$.

These provide an inertial mass term to the effective action of each the forward and backward paths.

The remaining terms, ignoring Bessel function derivatives as before,

$$\begin{aligned}\Phi_s^{int} &= -\frac{Sq^2\pi}{2a^2} \int dk k \int_0^T dt \int_0^t ds \frac{\cos ckQ(t-s)}{ckQ^4r_v^2} \ddot{\mathbf{X}}(s) \cdot \dot{\mathbf{X}}(t) \\ &\quad J_0(k|\mathbf{X}(t) - \mathbf{X}(s)|) + \text{etc.} \\ &\approx -\frac{2Sq^2\pi^2r_v^2}{2a^2c} \int_0^T dt \int_0^t ds \frac{\cos ckQ(t-s)}{ckQ^4r_v^2} \ddot{\mathbf{X}}(s) \cdot \dot{\mathbf{X}}(t) \\ &\quad J_0(k|\mathbf{X}(t) - \mathbf{X}(s)|) + \text{etc.}\end{aligned}\quad (4.74)$$

This is much smaller than the log divergent boundary term. Note that we've neglected the smaller still J_2 terms. By varying the $\ddot{X}\dot{X}$ terms with respect to X , we would obtain a small third order time derivative term, $\ddot{X}$, in the equations of motion. In an attempt to explain their numerical simulation results, Mertens et. al.^{41,41,69} artificially introduce a third order term by expanding the energy functional assuming both position and velocity as collective coordinates. This, of course, is a misapplication of the collective coordinate formalism, where a collective coordinate is meant to replace a continuous symmetry that the soliton breaks. The freedom they introduced by assuming velocity as a collective variable is not actually available in the original problem.

Consider the cosine term next. We can re-express this damping in terms of various damping functions

$$\begin{aligned}
\Phi_c = \int_0^T dt \int_0^t ds & \left(\gamma_{||}(t-s, |\mathbf{X}(t) - \mathbf{X}(s)|) \dot{\mathbf{X}}(s) \cdot \mathbf{X}(t) \right. \\
& + \gamma_{\Delta}(t-s, |\mathbf{X}(t) - \mathbf{X}(s)|) \dot{\mathbf{X}}(s) \hat{\mathbf{X}}(s) \cdot \hat{\mathbf{e}}_{\Delta}(\mathbf{X}(t) \cdot \hat{\mathbf{e}}_{\Delta}) \\
& \left. + \gamma_{\perp}(t-s, |\mathbf{X}(t) - \mathbf{X}(s)|) \dot{\mathbf{X}}(s) \hat{\mathbf{X}}(s) \cdot \hat{\mathbf{e}}_{\perp}(\mathbf{X}(t) \cdot \hat{\mathbf{e}}_{\perp}) \right) + \text{etc.}
\end{aligned} \tag{4.75}$$

where

$$\begin{aligned}
\gamma_{||}(t-s, \Delta) &= \frac{S^2 J \pi q^2}{2} \int dk k \frac{\cos \omega_0(t-s) J_0(k\Delta)}{Q^2 r_v^2} \\
\gamma_{\Delta}(t-s, \Delta) &= \frac{S^2 J \pi q^2}{2} \int dk k \frac{\cos \omega_0(t-s) J_2(k\Delta)}{Q^2 r_v^2} \\
\gamma_{\perp}(t-s, \Delta) &= -\gamma_{\Delta}(t-s, \Delta)
\end{aligned} \tag{4.76}$$

Note we cannot perform the k integrals in analytic form due to the $\omega_0 = ckQ$ argument in the cosine.

The damping forces depend on the previous motion of the vortex. These memory effects appear as averages over Bessel functions—this form is because the vortex exists in a 2D system. The first damping term is of the regular form, that is, a force acting in the opposite direction to the particle velocity. The next damping, γ_{Δ} , is the same as the first if the vortex travels in a straight line, however, for a curved path, is dependent on its change in direction. The last damping, $\gamma_{\perp}$, contributes damping perpendicular to the γ_{Δ} damping, which, in the case of a slowly curving path, is transverse to the vortex motion.

Comparing with the dissipation results of Slonczewski⁵⁹, although we find frequency dependent dissipation (via the $kX(t)$ coupling in the Bessel functions), we do not see any of the same small frequency behaviour predicted by Slonczewski. Likely, his treatment considers a different source of dissipation than the contribution considered here. As noted in section 4.2.2, this dissipation arises due to the same scattering processes that yield an inertial energy. In Slonczewski's treatment, on the other hand, his inertial energy calculation is for intermediate distance magnon scattering, while his dissipation arises from far field scattering.

4.3.5 Interpreting the real part

In the paper of Caldeira and Leggett⁵, the real part of the influence functional is interpreted as the correlation of forces in the classical regime. The real phase of their influence functional is

$$\Gamma = \sum_k \frac{C_k^2}{2m\omega_k} \coth \frac{\hbar\omega_k\beta}{2} \int_0^T dt \int_0^t ds (x(t) - y(t)) \cos \omega_k(t-s) (x(s) - y(s)) \quad (4.77)$$

which they compare to the contribution of a normally distributed classical fluctuating force $F(t)$ with correlation $\langle F(t)F(s) \rangle = A(t-s)$

$$\tilde{\Gamma} = \frac{1}{\hbar} \int_0^T dt \int_0^t ds (x(t) - y(t)) A(t-s) (x(s) - y(s)) \quad (4.78)$$

Since these terms have the same form, the real part of the influence functional must be interpretable as the correlation of forces in the classical regime.

The real phase of the vortex influence functional is, after substitution for f_k ,

$$\Gamma = \frac{Sq^2}{2a^2} \int d^2k \coth \frac{\hbar\omega_k\beta}{2} \int_0^T dt \int_0^t ds \left(\dot{\mathbf{X}}(t) e^{i\mathbf{k}\cdot\mathbf{X}(t)} - \dot{\mathbf{Y}}(t) e^{i\mathbf{k}\cdot\mathbf{Y}(t)} \right) \cdot \hat{\varphi}_{\mathbf{k}} \\ \frac{\cos \omega_k(t-s)}{kQ^3r_v^2} \left(\dot{\mathbf{X}}(s) e^{-i\mathbf{k}\cdot\mathbf{X}(s)} - \dot{\mathbf{Y}}(s) e^{-i\mathbf{k}\cdot\mathbf{Y}(s)} \right) \cdot \hat{\varphi}_{\mathbf{k}} \quad (4.79)$$

The integral over φ_k can be performed exactly as was done for the imaginary part yielding Bessel function pre-factors

$$\Gamma = \frac{\pi Sq^2}{2a^2} \int dk k \coth \frac{\hbar\omega_k\beta}{2} \int_0^T dt \int_0^t ds \frac{\cos \omega_k(t-s)}{kQ^3r_v^2} \\ \left(\dot{\mathbf{X}}(t) \cdot \dot{\mathbf{X}}(s) J_0(k|\mathbf{X}(t) - \mathbf{X}(s)|) + \dot{\mathbf{X}}(t) \dot{\mathbf{X}}(s) \left((\hat{\mathbf{X}}(t) \cdot \hat{\mathbf{e}}_{\Delta})(\hat{\mathbf{X}}(s) \cdot \hat{\mathbf{e}}_{\Delta}) \right. \right. \\ \left. \left. - (\hat{\mathbf{X}}(t) \times \hat{\mathbf{e}}_{\Delta}) \cdot (\hat{\mathbf{X}}(s) \times \hat{\mathbf{e}}_{\Delta}) \right) J_2(k|\mathbf{X}(t) - \mathbf{X}(s)|) \right) + \text{etc.} \quad (4.80)$$

Integrating by parts twice to cast this into a similar form to the regular dissipation term of Caldeira and Leggett

$$\Gamma = \frac{\pi Sq^2}{2a^2} \int dk \omega_k^2 \coth \frac{\hbar\omega_k\beta}{2} \int_0^T dt \int_0^t ds \frac{\cos \omega_k(t-s)}{Q^3r_v^2} \\ \left(\mathbf{X}(t) \cdot \mathbf{X}(s) J_0(k|\mathbf{X}(t) - \mathbf{X}(s)|) + ((\mathbf{X}(t) \cdot \hat{\mathbf{e}}_{\Delta})(\mathbf{X}(s) \cdot \hat{\mathbf{e}}_{\Delta}) \right. \\ \left. - (\mathbf{X}(t) \times \hat{\mathbf{e}}_{\Delta}) \cdot (\mathbf{X}(s) \times \hat{\mathbf{e}}_{\Delta})) J_2(k|\mathbf{X}(t) - \mathbf{X}(s)|) \right) + \text{etc.} \quad (4.81)$$

where, as usual, we neglect derivatives of the Bessel functions since their derivatives provide higher order corrections. There are additional boundary terms depending only on initial and final positions that don't affect the vortex dynamics.

The real phase can be interpreted as the correlation of forces. However, here the fluctuating forces are now vector forces and there are correlations between various components of the fluctuating forces. The appearance of the various Bessel functions, arising because the vortex is an extended object in 2D, differs from the treatment of Caldeira and Leggett because of a different density of states of the environmental modes.

4.4 Discussion of vortex effective dynamics

Markovian approximation

We can apply the Markovian approximation as in Castro Neto and Caldeira's treatment of solitons⁴². That is, approximate $\gamma(t) \rightarrow \gamma \frac{\delta(t)}{\omega_0}$ and similarly in the force correlation integral, (4.81), giving

$$\Gamma_M = \frac{\tilde{A}(\beta)}{\hbar} \int_0^T dt (\mathbf{X}(t) - \mathbf{Y}(t)) \cdot (\mathbf{X}(t) - \mathbf{Y}(t)) \quad (4.82)$$

where

$$\tilde{A}(\beta) = S^2 J \pi q^2 \hbar \int dk k \frac{1}{Q^2 r_v^2} \coth \frac{\hbar \omega_k \beta}{2} \quad (4.83)$$

Note, all J_2 terms disappear in this approximation.

In this limit, the longitudinal damping coefficient, (4.75) becomes

$$\begin{aligned} \eta &= \frac{S \pi q^2}{a^2} \int dk \frac{1}{Q^3 r_v^2} \\ &= \frac{S \pi q^2}{a^2} \end{aligned} \quad (4.84)$$

The classical fluctuation-dissipation theorem is now satisfied in the high temperature limit ($\coth x \rightarrow \frac{1}{x}$)

$$\tilde{A}(\beta) = 2k_B T \eta \quad (4.85)$$

where T here denotes temperature.

This limit corresponds to the limit where the timescale of interest is much greater than the correlation time of the magnons.

4.4.1 Comparison with radiative dissipation

The dissipation found in the Markovian approximation can be compared with the over-simplified calculation performed using second order perturbation theory. There, assuming only the emission of a magnon and no inter-magnon scattering, we found that the dissipation rate was given by the integral

$$\gamma = 2\pi \int d^2k \frac{Sq^2}{2a^2kQ^3r_v^2} \left(\dot{\mathbf{X}} \cdot \hat{\chi}_k \right)^2 \delta(\hbar ckQ) \quad (4.86)$$

where we evaluated this integral in section 4.2.2. Comparing the k dependence of this integral with that of the damping coefficient in equation (4.84), we find they differ only by the δ -function. The dissipation is now dependent on the entire magnon spectrum. In the previous calculation, we made the simplifying assumption that there were initially no magnons and hence only zero energy magnons could be scattered.

In the Markovian limit, the effective damping force found in the imaginary part of the influence functional phase gives roughly the energy dissipation

$$\begin{aligned} E_{diss} &\sim \int dX \cdot \eta \dot{\mathbf{X}} \\ &= \int dt \eta \dot{X}^2 \\ &= \int dt \frac{S\pi q^2}{a^2} \dot{X}^2 \\ &= \frac{S\pi q^2 T}{a^2} \dot{X}^2 \end{aligned} \quad (4.87)$$

Here, the full spectrum of magnons is excited, with probability of finding a certain k state weighted by its corresponding Boltzmann factor. Thus, even assuming no vortex inertial energy, we can find scattering between infinitesimally spaced k states throughout the spectrum.

4.4.2 Extending results to many vortices

The entire treatment can be repeated for a collection of vortices. Assuming the vortices are well enough separated to neglect core interactions, the unperturbed

spin configuration is

$$\begin{aligned}\phi_{tot} &= \sum_{i=1}^n q_i \chi(\mathbf{X}_i) \\ \theta_{tot} &= \sum_{i=1}^n \theta_v(\mathbf{r} - \mathbf{X}_i)\end{aligned}\tag{4.88}$$

where $\mathbf{X}_i$ denotes the center of the i th vortex. The center coordinates are elevated to operators within the collective coordinate formalism. Expanding the Lagrangian in terms of this spin field, without magnon interactions, we find gyrotropic momentum terms and inter-vortex potentials

$$\mathcal{L}_v^0 = \sum_i \left(-E_{v,i} + \mathbf{P}_{gyro,i} \cdot \dot{\mathbf{X}}_i + 2S^2 J \pi \sum_{j \neq i} q_i q_j \ln \frac{X_{ij}}{r_v} \right) \tag{4.89}$$

where $E_{v,i}$ is the unimportant rest energy of the vortices, $\mathbf{P}_{gyro,i} = -\frac{\pi S q_i p_i}{a^2} \mathbf{X}_i \times \hat{\mathbf{z}}$ is the vector potential giving the gyrotropic force, and the last term accounts for inter-vortex interactions.

The magnon interactions are treated to leading order only—the zero-point energy shifts do not affecting dynamics, and higher order dissipation is not treated here. There is a one magnon coupling with the vortex velocities $\dot{\mathbf{X}}_i$ to the magnons that is integrated over in the influence functional.

The resulting influence functional now has effective action terms coupling not only the forward and backward paths of the same vortex, but also the paths for different vortices. Without going through all the details, the general results are presented. Were we to neglect all inter-vortex terms, the influence functional would simply be the product over each single vortex influence functional.

Including inter-vortex terms to leading order now, the mass tensor is exactly the same as that found using second order perturbation theory (see section 4.2.1) and, in fact, the calculations here are nearly identical to those. The mass tensor is

$$\begin{aligned}M_{ij} &= \int d^2k \frac{q_i q_j}{J a^4 k^2 Q^4 r_v^2} \left(\hat{\mathbf{X}}_i \cdot \hat{\chi}_k \right) \left(\hat{\mathbf{X}}_j \cdot \hat{\chi}_k \right) e^{i\mathbf{k} \cdot (\mathbf{X}_i - \mathbf{X}_j)} \\ &= \frac{\pi q_i q_j r_v^2}{J a^4} \begin{cases} \ln \frac{R_S}{r_{ij}} + \frac{1}{2} \left((\hat{\mathbf{X}}_i \cdot \hat{\mathbf{e}}_{ij})(\hat{\mathbf{X}}_j \cdot \hat{\mathbf{e}}_{ij}) \right. \\ \left. - (\hat{\mathbf{X}}_i \times \hat{\mathbf{e}}_{ij}) \cdot (\hat{\mathbf{X}}_j \times \hat{\mathbf{e}}_{ij}) \right), & i \neq j; \\ \ln \frac{R_S}{\sqrt{a^2 + r_v^2}}, & i = j. \end{cases} \end{aligned} \tag{4.90}$$

where $\hat{\mathbf{e}}_{ij} = \frac{\mathbf{X}_i - \mathbf{X}_j}{|\mathbf{X}_i - \mathbf{X}_j|}$.

There are inter-vortex damping forces behaving essentially like the single vortex damping forces: there exist forces longitudinal and transverse to the motion of

a vortex, however, acting on a second vortex. The damping decreases as a function of vortex separation as $\sim J_0(kr_{ij})$. This dissipation is thus quite small when we assume well separated vortices, in keeping with previous calculations (refer to the inter-vortex forces calculation in section 3.1).

Similarly, in the force correlation integral, we find that the fluctuating forces acting on various vortices are inter-correlated. This shouldn't be surprising at all: we have damping terms intermingling the motion of vortex pairs so that we should therefore expect that the fluctuating forces on these vortices are inter-dependent.

The final effective density matrix propagator becomes

$$\begin{aligned}
J(\mathbf{X}_i, \mathbf{Y}_i; \mathbf{X}'_i, \mathbf{Y}'_i) &= \prod_i \int_{\mathbf{X}_i, \mathbf{Y}_i}^{\mathbf{X}'_i, \mathbf{Y}'_i} \mathcal{D}[\mathbf{X}_i(t), \mathbf{Y}_i(t)] \exp \frac{i}{\hbar} (\mathcal{S}_v[\mathbf{X}_i(t)] - \mathcal{S}_v[\mathbf{Y}_i(t)]) \\
&\exp -\frac{1}{\hbar^2} \sum_i \int_0^T \int_0^t dt ds \left(\sum_{i,j} A_{ij}(t-s) \mathbf{X}_i(t) \cdot \mathbf{X}_j(s) J_0(k|\mathbf{X}_i(t) - \mathbf{X}_j(s)|) \right. \\
&+ \sum_i A_{ii}(t-s) \left((\mathbf{X}_i(t) \cdot \hat{\mathbf{e}}_{\Delta_i})(\mathbf{X}_i(s) \cdot \hat{\mathbf{e}}_{\Delta_i}) \right. \\
&\left. \left. - (\mathbf{X}_i(t) \times \hat{\mathbf{e}}_{\Delta_i}) \cdot (\mathbf{X}_i(s) \times \hat{\mathbf{e}}_{\Delta_i}) \right) J_2(k|\mathbf{X}_i(t) - \mathbf{X}_i(s)|) \right) + \text{etc.} \quad (4.91)
\end{aligned}$$

where the force correlations as applied to vortices i and j are

$$A_{ij}(t-s) = \frac{\hbar \pi S q_i q_j}{2a^2} \int dk \frac{\coth \frac{\hbar \omega_k \beta}{2}}{Q^3 r_v^2} \omega_k^2 \cos \omega_k(t-s) \quad (4.92)$$

The vortex effective action has been redefined to include the inertial mass and damping terms

$$\begin{aligned}
\mathcal{S}_v &= \int dt \left(\mathcal{L}_v^0 + \sum_{i,j} \frac{1}{2} M_{ij} \dot{\mathbf{X}}_i \cdot \dot{\mathbf{X}}_j \right. \\
&- \int_0^t ds \left(\sum_{i,j} \gamma_{||}^{ij}(t-s, |\mathbf{X}_j(t) - \mathbf{X}_i(s)|) \dot{\mathbf{X}}_i(s) \cdot \mathbf{X}_j(t) \right. \\
&+ \sum_i \gamma_{\Delta}^i(t-s, |\mathbf{X}_i(t) - \mathbf{X}_i(s)|) \dot{\mathbf{X}}_i(s) (\hat{\mathbf{X}}_i(s) \cdot \hat{\mathbf{e}}_{\Delta_i}) (\mathbf{X}_i(t) \cdot \hat{\mathbf{e}}_{\Delta_i}) \\
&\left. \left. + \gamma_{\perp}^i(t-s, |\mathbf{X}_i(t) - \mathbf{X}_i(s)|) \dot{\mathbf{X}}_i(s) (\hat{\mathbf{X}}_i(s) \cdot \hat{\mathbf{e}}_{\perp_i}) (\mathbf{X}_i(t) \cdot \hat{\mathbf{e}}_{\perp_i}) \right) \right) \quad (4.93)
\end{aligned}$$

where the damping functions are from (4.76)

$$\begin{aligned}\gamma_{||}^{ij}(t-s, \Delta) &= \frac{S^2 J \pi q_i q_j}{2} \int dk k \frac{\cos \omega_0(t-s) J_0(k\Delta)}{Q^2 r_v^2} \\ \gamma_{\Delta}^i(t-s, \Delta) &= \frac{S^2 J \pi q_i^2}{2} \int dk k \frac{\cos \omega_0(t-s) J_2(k\Delta)}{Q^2 r_v^2} \\ \gamma_{\perp}^i(t-s, \Delta) &= -\gamma_{\Delta}^i(t-s, \Delta)\end{aligned}\tag{4.94}$$

Note in the limit of slow motion and large inter-vortex separation, that is, $J_0 \rightarrow 1$ for same vortex terms and all others are negligible, this effective action has the same form for each vortex as found in the quantum Brownian motion described by Caldeira and Leggett⁵, however, with inter-vortex terms introducing Coulomb-like forces.

4.4.3 Frequency dependent motion

Perhaps a better way of understanding the role of the Bessel function pre-factors is to decompose them according to the sum rules

$$J_\nu(k|\mathbf{x}-\mathbf{y}|) = \sum_{m=-\infty}^{\infty} J_m(kx) J_{\nu+m}(ky) e^{i(\nu+m)(\phi_x - \phi_y)}\tag{4.95}$$

Denote $\mathbf{X}_{km}^i = J_m(kX_i) e^{im\phi_i} \mathbf{X}$. The effective Lagrangian is transformed to

$$\begin{aligned}\mathcal{L}_v = \mathcal{L}_v^0 &+ \sum_m \int dk \frac{1}{2} \sum_{i,j} M_k^{ij} \dot{\mathbf{X}}_{km}^i \cdot \dot{\mathbf{X}}_{km}^j + \frac{1}{2} \sum_i M_k^{ii} \dot{\mathbf{X}}_{km}^i \cdot \dot{\mathbf{X}}_{k,m+2}^i e^{i2\phi_i} \\ &- \int_0^t ds \left(\sum_{i,j} \gamma_{||k}^{ij}(t-s) \dot{\mathbf{X}}_{km}^i(s) \cdot \mathbf{X}_{km}^j(t) \right. \\ &+ \sum_i \gamma_{\Delta k}^i(t-s) e^{i2\phi_{is}} \dot{X}_{km}^i(s) (\hat{\mathbf{X}}^i(s) \cdot \hat{\mathbf{e}}_{\Delta_i}) (\mathbf{X}_{k,m+2}^i(t) \cdot \hat{\mathbf{e}}_{\Delta_i}) \\ &\left. + \gamma_{\perp k}^i(t-s) e^{i2\phi_{is}} \dot{X}_{km}^i(s) (\hat{\mathbf{X}}^i(s) \cdot \hat{\mathbf{e}}_{\perp_i}) (\mathbf{X}_{k,m+2}^i(t) \cdot \hat{\mathbf{e}}_{\perp_i}) \right)\end{aligned}\tag{4.96}$$

where $M_k^{ij} = \frac{\pi q_i q_j}{J a^4 k Q^4 r_v^2}$, recalling that the mass tensor can be expressed as an integral over k with Bessel function factors (refer to section 4.2.1). The new damping function is defined as

$$\begin{aligned}\gamma_{||k}^{ij}(t-s) &= \frac{S^2 J \pi q_i q_j}{2} \frac{k}{Q^2 r_v^2} \cos \omega_0(t-s) \\ \gamma_{\Delta k}^i(t-s) &= \frac{S^2 J \pi q_i^2}{2} \frac{k}{Q^2 r_v^2} \cos \omega_0(t-s) \\ \gamma_{\perp k}^i(t-s) &= -\gamma_{\Delta k}^i(t-s, \Delta)\end{aligned}\tag{4.97}$$

The real part of the influence functional can be re-expressed now as

$$\begin{aligned} \Gamma = & \frac{1}{\hbar} \sum_{ijm} \int dk \int_0^T dt \int_0^t ds A_k^{ij}(t-s) (\mathbf{X}_{km}^i(t) - \mathbf{Y}_{km}^i(t)) \cdot (\mathbf{X}_{km}^j(s) - \mathbf{Y}_{km}^j(s)) \\ & + \delta_{ij} A_k^{ii}(t-s) e^{i2\phi_{it}} \mathbf{X}_{km}^i(t) \cdot \left(\hat{\mathbf{e}}_{\Delta_i} (\mathbf{X}_{k,m+2}^i(s) \cdot \hat{\mathbf{e}}_{\Delta_i}) \right. \\ & \left. - \hat{\mathbf{e}}_{\perp_i} \cdot (\mathbf{X}_{k,m+2}^i(s) \cdot \hat{\mathbf{e}}_{\perp_i}) \right) + \text{etc. in } XY \text{ and } Y^2 \end{aligned} \quad (4.98)$$

where

$$A_k^{ij}(t-s) = \frac{\hbar \pi S q_i q_j}{2a^2} \frac{\coth \frac{\hbar \omega_k \beta}{2}}{Q^3 r_v^2} \omega_k^2 \cos \omega_k(t-s) \quad (4.99)$$

Recall that the density matrix propagator is not simply the product of non-interacting forward and backward paths. As in (4.91), we also have damping terms coupling the forward and backward paths.

Thus, we find that the motion of the collection of vortices behaves as interacting Brownian particles; however, with frequency dependent damping and fluctuating forces. The formalism of Caldeira and Leggett⁵ can be applied to each frequency component, with the added complexity of inter-vortex forces.

4.4.4 Summary

A collection of vortices are quantized by considering the small perturbations about them. This amounts to including vortex-magnons interactions. We considered two couplings in depth: a first order coupling between the vortex velocity and the magnon spin field, and a second order magnon coupling. All vortex-magnon couplings create dissipation via magnon radiative processes. We considered only the dissipation due to the first order coupling, first in perturbation theory and later via the influence functional. Higher order couplings also create dissipation, and may, in fact, contribute more significantly^{12,62}, however, these weren't considered here.

The one magnon coupling creates an inertial energy endowing the vortex with an effective mass. A collection of vortices are strongly coupled: in addition to the usual inter-vortex forces, there are inter-vortex inertial terms such as $\frac{1}{2} M_{ij} \dot{\mathbf{X}}_i \cdot \dot{\mathbf{X}}_j$ that are non-negligible. The zero point energy shift from the two magnon coupling is log divergent and, being due to the presence of the vortex, is considered the quantized vortex's zero point energy. Note, we did not calculate the full effect of this two magnon coupling, only that portion independent of magnon populations. This shift was calculated first by considering magnon scattering in Chapter 3 and next in this chapter by simply rewriting the interaction in terms of magnon creation/annihilation operators.

The influence functional reconfirms the effective mass calculations and gives explicitly the damping forces and corresponding fluctuating forces responsible for dissipation. These act longitudinally and transverse to the vortex motion. Again, a collection of vortices are coupled via the damping forces: damping forces due to the motion of a first vortex act on a second vortex. Damping forces depend on the entire history of the vortex dynamics.

Chapter 5

Conclusions

We study the dynamics of a collection of magnetic vortices in an easy plane two dimensional insulating ferromagnet. The system is approximated by a continuous spin field because we are only interested in the low energy response. The vortices interact with magnons via a variety of couplings. The effective dynamics bear many similarities to that in the more complex superfluid and superconducting vortex bearing systems.

We reviewed the derivations of the gyrotropic force and the inter-vortex force by expanding the vortex action about a stationary superposition of vortex solutions. We reviewed the inertial mass derivation by calculating vortex profile distortions when in motion and showed the equivalence of this method with ordinary perturbation theory. We reviewed magnon phase shift calculations and how these phase shifts modify the vortex zero point energy. By rewriting the scattering potential in terms of magnon creation and annihilation operators, we found an equivalence of the phase shift calculations with the immediate energy shift revealed in the second quantized form.

We suggest a new interpretation of the gyrotropic force as a Lorentz-type force with the vortex vorticity behaving like charge, $4\pi\epsilon_0 q$ (in SI units), in an effective perpendicular magnetic field, $\mathbf{B} = \frac{S^2 J}{4\epsilon_0 r_v} p_i \hat{z}$, due to the vortex's own out-of-plane spins. We rewrite the effective action term giving the gyrotropic force instead as a vector potential shift in the vortex momentum. This momentum term was then verified by direct integration of the operator generating translations. The vector potential possesses gauge freedom, allowed by the same freedom of gauge in the Berry's phase.

We next employed the Feynman-Vernon influence functional formalism, assuming the vortex-magnon systems are initially uncoupled with the magnons in thermal equilibrium (thus introducing temperature). The systems interact and entangle. The dynamics of the vortices were isolated by tracing over magnons. The resulting effective vortex motion is acted upon by longitudinal and transverse damping forces. Before now, no damping force acting transverse to the vortex motion has been suggested in a magnetic system.

The vortex is a stable solution of the easy plane ferromagnet. As such, when we expand about it to quantize magnons in its presence, we find no linear coupling between the two fields. However, the vortex is a stationary solution, so that

setting it into motion, we find a first order coupling between the magnon field and the vortex velocity.

This lowest order coupling, responsible for endowing the vortex with an effective mass, is dissipative and yields effective damping forces acting on a moving vortex. The damping forces are accompanied by fluctuating forces that average to zero and with time correlations such that the fluctuation-dissipation theorem is satisfied in a generalized way.

We found both longitudinal and transverse damping forces dependent on the prior motion of the vortex. For a collection of vortices, the damping forces also act between vortices: the motion of a first vortex causes a damping force to act on a second vortex. Correspondingly, there are non-zero correlations between forces acting on two different vortices.

The vortex dynamics were described by the propagator of the vortex reduced density matrix. The forward and backward paths are coupled, as already described for quantum Brownian motion by Caldeira and Leggett⁵. The damping forces possess memory effects, a common feature in general when describing a soliton as a quantum Brownian particle⁴². In our two dimensional system, however, we found additional Bessel function factors. These considerably complicate the extraction of a spectral function describing the ensuing Brownian motion. By decomposing the vortex motion in a basis of Bessel functions, we find that the various frequency components behave as a coupled ensemble of quantum Brownian particles.

5.1 Open questions

The analogy of a vortex as a charged particle in a magnetic field can be extended. For instance, there should be excitations within the gauge field giving the gyrotropic momentum. The magnetic field is a result of the out-of-plane spins at the vortex center. Perhaps, gauge fluctuations are related to vortex core flips.

Future work on magnetic vortex motion should check the relative importance of higher order dissipative couplings. The basic motion of a small collection of vortices can be examined now including inertial and damping forces. For instance, one could verify the claim of Slonczewski⁵⁹ that damping forces acting on a vortex pair only decay circular orbits inward and parallel ones outward. The similarities with superfluid vortices should be further examined by attempting to calculate the influence functional of a superfluid vortex and, likewise, the Aharonov-Bohm interference effects of magnons passing a moving vortex.

Appendix A

Some mechanics

A classical system is describable by its Lagrangian, which is a function of the system coordinates q_i and velocities $\dot{q}_i$

$$\mathcal{L}(q_i, \dot{q}_i, t)$$

The action of the system is defined by

$$\mathcal{S}[q_i(t)] = \int_0^T dt L(q_i, \dot{q}_i, t) \quad (\text{A.1})$$

The equations of motion of the system are given by the principle of least action, otherwise known as Hamilton's principle, stating that the system evolves from initial state $\{q_i(0)\}$ to final state $\{q_i(T)\}$ via the path $q_i(t)$ that extremizes the action, $\mathcal{S}$.

Given that $\mathcal{L} = \mathcal{L}(q_i, \dot{q}_i, t)$, extremizing the action we arrive at the Euler-Lagrange equations

$$\frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{q}_i} - \frac{\partial \mathcal{L}}{\partial q_i} = 0 \quad (\text{A.2})$$

Alternatively, we can describe the system by its Hamiltonian. We transform from the Lagrangian to the Hamilton via a Legendre transformation

$$H(q_i, p_i, t) = \sum_i p_i \dot{q}_i - \mathcal{L}(q_i, \dot{q}_i, t) \quad (\text{A.3})$$

where we've defined the conjugate momenta p_i defined by

$$p_i = \frac{\partial \mathcal{L}}{\partial \dot{q}_i} \quad (\text{A.4})$$

Hamilton's equations are a restatement of (A.2) and (A.4)

$$\frac{\partial q_i}{\partial t} = \frac{\partial H}{\partial p_i}; \quad \frac{\partial p_i}{\partial t} = -\frac{\partial H}{\partial q_i} \quad (\text{A.5})$$

For example, consider a particle of mass m , position x , residing in a potential $V(x)$. The Hamiltonian is simply the total energy of the system

$$H(x, p, t) = \frac{p^2}{2m} + V(x) \quad (\text{A.6})$$

where the conjugate momentum $p = m\dot{x}$, as usual. The Lagrangian is then given as the difference in kinetic and potential energies

$$\mathcal{L}(x, \dot{x}, t) = \frac{1}{2}m\dot{x}^2 - V(x) \quad (\text{A.7})$$

Application of either the Euler-Lagrange equation or Hamilton's equations yields Newton's second law of motion, $F = m\ddot{x}$, where the force $F = -\frac{\partial}{\partial x}V(x)$.

Define the Poisson bracket $\{\cdot, \cdot\}_{q,p}$ for a system with coordinate q and conjugate momentum p via

$$\{A, B\}_{q,p} = \frac{\partial A}{\partial q} \frac{\partial B}{\partial p} - \frac{\partial A}{\partial p} \frac{\partial B}{\partial q} \quad (\text{A.8})$$

Note that $\{q, p\}_{q,p} = 1$. The above can be easily generalized to a field theory by substituting $q \rightarrow \tilde{\phi}(x)$ and $p \rightarrow \tilde{\pi}(x)$ and replacing all simple derivatives by functional derivatives.

Going over to quantum mechanics, to quantize the motion of the system, we impose the commutation relations

$$[q, p] = i\hbar \quad (\text{A.9})$$

In 1925, P. A. M. Dirac¹⁰ observed that proper quantum mechanical relations followed under the substitution

$$\{\cdot, \cdot\}_{q,p} \rightarrow \frac{1}{i\hbar}[\cdot, \cdot]$$

In a spin system, using coordinate ϕ and conjugate momentum $S \cos \theta$ we can verify directly the classical version of $[S_i, S_j] = i\hbar \varepsilon_{ijk} S_k$, that is,

$$\{S_i, S_j\}_{\phi, S \cos \theta} = \varepsilon_{ijk} S_k \quad (\text{A.10})$$

where we define $S = S(\sin \theta \cos \phi, \sin \theta \sin \phi, \cos \theta)$. However, spin being an essentially quantum concept, we must bear in mind that when speaking of spin directions given by (ϕ, θ) , we mean the spin state of highest probability to be found in that direction.

A.1 Imaginary time path integral

Consider a system in thermal equilibrium at temperature τ . If we decompose the system Hamiltonian into a set of eigenstates $\xi_n(x)$ with eigenenergies E_n , then the probability of observing the system in eigenstate n is proportional to $e^{-\frac{E_n}{k_B\tau}}$ where k_B is Boltzmann's constant¹⁵. The density matrix for this system is

$$\rho(x', x) = \sum_n \xi_n(x') \xi_n^*(x) e^{-\beta E_n} \quad (\text{A.11})$$

where $\beta = (k_B\tau)^{-1}$.

Compare this with the quantum propagator decomposed into this same basis:

$$\begin{aligned} K(x', T; x, 0) &= \langle x' | \exp -\frac{iHT}{\hbar} | x \rangle \\ &= \sum_n \langle x' | \xi_n \rangle \exp -\frac{iE_n T}{\hbar} \langle \xi_n | x \rangle \\ &= \sum_n \xi_n(x') \xi_n^*(x) \exp -\frac{iE_n T}{\hbar} \end{aligned} \quad (\text{A.12})$$

We see that the density matrix is formally identical to the propagator corresponding to an imaginary time interval $T = -i\beta\hbar$. In fact, if we consider the equation of motion of the density matrix found by taking the derivative of (A.11) with respect to β ¹⁶

$$-\frac{\partial \rho}{\partial \beta} = \sum_n E_n \xi_n(x') \xi_n^*(x) e^{-\beta E_n} \quad (\text{A.13})$$

Recall that $E_n \xi_n(x') = H \xi_n(x')$. If we understand $H_{x'}$ to act only on x' , we can write

$$-\frac{\partial \rho(x', x)}{\partial \beta} = H_{x'} \rho(x', x) \quad (\text{A.14})$$

We know how to evaluate the propagator as a path integral for simple Hamiltonians involving only the system coordinates and their conjugate momenta. For example, for the Hamiltonian

$$H = -\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + V(x) \quad (\text{A.15})$$

the solution over an infinitesimal time period ϵ is

$$K(x', \epsilon; x, 0) = \sqrt{\frac{m}{2\pi i \hbar \epsilon}} \exp \frac{i}{\hbar} \left(\frac{m}{2} \frac{(x' - x)^2}{\epsilon} - \epsilon V \left(\frac{x' + x}{2} \right) \right) \quad (\text{A.16})$$

which can be verified by direct substitution into

$$-\frac{\hbar}{i} \frac{\partial K(x', T; x, 0)}{\partial T} = H_{x'} K(x', T; x, 0) \quad (\text{A.17})$$

Now, under an infinitesimal interval in the density matrix $i\epsilon$ the solution is given by $\epsilon \rightarrow -i\epsilon$

$$\rho(x', x; \beta = \epsilon/\hbar) = \sqrt{\frac{m}{2\pi \hbar \epsilon}} \exp -\frac{1}{\hbar} \left(\frac{m}{2} \frac{(x' - x)^2}{\epsilon} + \epsilon V \left(\frac{x' + x}{2} \right) \right) \quad (\text{A.18})$$

which can be verified by direct substitution into (A.14).

Stringing many of these solutions together for successive intervals of time according to

$$\rho(x', x; \beta') = \int dx'' \rho(x', x'', \beta') \rho(x'', x; \beta) \quad (\text{A.19})$$

for intermediate x'' at β , we obtain a path integral formulation of the density matrix which is simply an imaginary time version of the propagator path integral, that is, with the substitution $T \rightarrow -i\beta\hbar$.

Appendix B

Quantization of classical solutions

Suppose we have a particle described by position x residing in a potential $V(x)$. Classically, the particle follows a path $x(t)$ that satisfies Newton's second law of motion. In quantum theory, the particle is no longer described by its position x , but by its wavefunction $\psi(x)$ giving a probability distribution of finding the particle at position x . If the energy is conserved, the wavefunction can be decomposed into energy eigenstates, $\psi_n(x)$, obeying Schroedinger's equation

$$H\psi_n = E_n\psi_n(x) \quad (\text{B.1})$$

where H is the Hamiltonian of the system, quantized by elevating the position and momentum variables to operators.

As preparation for a description of the quantization of a soliton solution, consider some of the finer points of quantization of classical particle solutions.

For the potential shown in Figure B.1, there are three extrema and hence three stationary classical solutions. The absolute minimum, $x = a$ is the classical ground state, having the lowest attainable energy.

In quantum mechanics, according to the uncertainty principle, a solution is not allowed to have zero momentum and a fixed position. Thus, even in its ground




Figure B.1: An illustrative potential of a one dimensional particle.

state there are fluctuations. Expanding $V(x)$ in a Taylor series, to lowest order the potential is harmonic about the minimum and we have simple harmonic excitations with frequency $\omega^2 = V''(x = a)$ and energies

$$E_n = V(a) + (n + \frac{1}{2})\hbar\omega \quad (\text{B.2})$$

The ground state energy becomes $E_0 = V(a) + \frac{1}{2}\hbar\omega$. The additional $\frac{1}{2}\hbar\omega$ is the zero-point energy due to quantum fluctuations.

The solution $x = c$ is a second stable classical solution. Quantum mechanically, there are again fluctuations about this solution that give a similar excitation spectrum. In this case, however, since this is an excited state, there are possibly tunneling processes that relax the state to its ground state about $x = a$. In a field theory, this stable excited state is the analogue of a soliton solution (while the tunneling processes are analogous to instantons). However, for the magnetic solitonic solutions considered here, these excited states belong to separate topological sectors of the solution space so that there is effectively an infinite energy barrier to the ground state.

The classical solution $x = b$ is unstable and would thus correspond to an imaginary frequency. There are hence no set of quantum levels formed about it.

Another interesting analogy to consider is the case of a constant potential, $V(x) = V$. In that case, there is no clear choice of minimum about which to expand and, should we attempt to, we would find everywhere $\omega = 0$. Of course, in quantum mechanics, the proper solutions to consider are the plane waves e^{ikx} with energies

$$E_n = V + \frac{1}{2m}(\hbar k_n)^2 \quad (\text{B.3})$$

where $\hbar k_n = p_n$ are the momenta of these states. In field theory, we find a zero frequency mode, or Goldstone mode, for every broken continuous symmetry. Further, for each of these broken symmetries, we find a corresponding conserved momentum, analogous to the conserved p_n in the particle case.

B.1 Quantizing soliton solutions

In field theory, quantizing a soliton follows analogously to the regular quantization of a classical solution. The language is changed somewhat however. For instance, the ground state of the particle, $x = a$, is quantized to a hierarchy of simple harmonic excitations. In field theory, we call the absolute potential minimum the ground state, or the vacuum state. The hierarchy of perturbative excitations are interpreted as mesons or quasiparticles. In our system, these are the magnons.

When we expand about the solitonic excited state (analogous to the second minimum, $x = c$), generally the quasiparticles are modified by the soliton presence. In the simple particle case, this corresponds to the general case where $V''(a) \neq V''(c)$. In the particle case, the hierarchy of simple harmonic states are interpreted as excited states about the minima.

In a field theory, the quasiparticles are generally extended states and, in the presence of a soliton, are shifted but still extended. In some cases, the soliton can trap a few quasiparticle modes. These bound modes are interpreted as soliton excited states. The remaining, extended states are interpreted as unshifted quasiparticles, while all energy shifts due to the soliton are attributed to the zero-point energy of the quantized soliton.

The soliton acts perturbatively on the extended states, it itself being localized in space, as a scattering center. Asymptotically far from the soliton center, the quasiparticles are simply phase shifted. Suppose that the relative phase shift between the incoming and outgoing waves is $\delta(k)$, a function of the wavevector k .

By enforcing periodic boundary conditions[†] on both the unperturbed wavevector k and the scattered wavevector q

$$\begin{aligned} Lk_n &= 2n\pi \\ Lq_n - \delta(q_n) &= 2n\pi \end{aligned} \tag{B.4}$$

we fix the allowed k and q values. In the $L \rightarrow \infty$ limit, these allowed values merge to a continuum and the sum over k -states is replaced by an integral

$$\sum_k \rightarrow \frac{L}{2\pi} \int_{-\infty}^{\infty} dk$$

The energy correction to the soliton solution, taken as the modification to the zero point energy of the vacuum, is thus, noting that $\omega(q) = \omega(k + \frac{\delta}{L})$,

$$\begin{aligned} \Delta E &= \frac{1}{2} \hbar \sum_k \omega(q) - \omega(k) \\ &= \frac{1}{2} \hbar \frac{L}{2\pi} \int_{-\infty}^{\infty} dk \frac{\partial \omega(k)}{\partial k} \frac{\delta(q)}{L} \\ &= \frac{1}{4\pi} \hbar \int_{-\infty}^{\infty} dk \frac{\partial \omega(k)}{\partial k} \delta(q) \end{aligned} \tag{B.5}$$

[†]Or alternatively, we could enforce fixed boundary conditions forcing k to be π -periodic rather than 2π -periodic. In that case, the asymptotic waveform must be modified from a plane wave to a cosine wavefunction and we find that the phase shift is also changed by a factor of 2. Thus, either set of boundary conditions is equivalent.

found by expanding to first order in δ .

In addition to small corrections to the quasiparticle continuum, the soliton might bind discrete levels in the quasiparticle spectrum. Those with $\omega = 0$ are due to a continuous symmetry broken by the soliton solution. These modes are dealt with using collective coordinates. There can also be $\omega \neq 0$ discrete modes. These are interpreted as soliton excited states. For an example of these, see the quantization of the quantum kink of the ϕ^4 theory⁵⁰—the magnetic vortex does not have any such excited states.

B.1.1 In a path integral formalism

Using path integrals, we can find the excitation spectrum of a system by taking the trace of the system's quantum propagator. We first review the simple case of a regular particle in an external potential and then generalize to field theory.

Semiclassical approximation for a single particle

The propagator of a single particle starting in position q_a at time 0 and ending in position q_b at time T is

$$K(q_b, T; q_a, 0) = \langle q_b | e^{-iHT/\hbar} | q_a \rangle \quad (\text{B.6})$$

where $H(q, p) = \frac{1}{2}p^2 + V(x)$, and, for simplicity, we've set $m = 1$.

Next, we set $q_a = q_b = q_0$ and integrate over the endpoint of the periodic orbit

$$\begin{aligned} G(T) &= \int dq_0 \langle q_0 | e^{-iHT/\hbar} | q_0 \rangle \\ &= \int dq_0 \sum_n \langle q_0 | \phi_n \rangle e^{-iE_n T/\hbar} \langle \phi_n | q_0 \rangle \\ &= \sum_n e^{-iE_n T/\hbar} \end{aligned} \quad (\text{B.7})$$

where $\{\phi_n\}$ denote a complete orthonormal set of eigenstates of H . This yields an expression giving the excitation spectrum of the Hamiltonian.

For a particle in a potential $V(x)$ with a minimum at $x = x_0$, the classical solution is simply $q_{cl} = x_0$. Expanding the potential in a power series about this solution

$$V(x) = V(q_{cl}) + V'(q_{cl})(q - q_{cl}) + \frac{1}{2}V''(q_{cl})(q - q_{cl})^2 + \mathcal{O}(\Delta x^3) \quad (\text{B.8})$$

the second term is zero since q_{cl} is a minimum of $V(x)$. The action expanded about this classical solution, $q(t) \rightarrow q_{cl} + q'(t)$, is now

$$\mathcal{S}[q(t)] = -V(x_0) + \int_0^T dt \frac{1}{2} \dot{q}'^2 - \frac{1}{2} \omega^2 q'^2 \quad (\text{B.9})$$

where $\omega^2 = V''(x_0)$, assumed positive (i.e. the classical solution is stable). Note, at this point, the boundary conditions of the periodic path are still not generally satisfied so that the new perturbed solution must now satisfy $q'(0) = q'(T) = q_0 - q_{cl}$.

The semiclassical approximation amounts to neglecting the $\mathcal{O}(\Delta x^3)$ and higher order terms. But the terms in q' are just the action of a simple harmonic oscillator. To evaluate the path integral

$$G_{SHO}(T) = \int dq_0 \int \mathcal{D}[q'(t)] e^{\frac{i}{\hbar} \mathcal{S}[q'(t)]} \quad (\text{B.10})$$

we expand again about the simple harmonic oscillator classical solution satisfying the appropriate boundary conditions. You may ask why we didn't immediately go from the beginning action and expand V in a Taylor series and approximate there. Although that would have proceeded identically, the additional step helps clarify what to do when expanding in a field theory admitting classical soliton solutions. The classical solution is now

$$q'_{cl} = A \cos \omega t + B \sin \omega t \quad (\text{B.11})$$

where the boundary conditions give

$$\begin{aligned} A &= q_0 - q_{cl} \\ A \cos \omega T + B \sin \omega T &= q_0 - q_{cl} \end{aligned} \quad (\text{B.12})$$

Evaluating this second classical contribution to the action gives

$$\mathcal{S}[q'_{cl}] = -2\omega(q_0 - q_{cl})^2 \frac{\sin^2 \omega T / 2}{\sin \omega T} \quad (\text{B.13})$$

The complete path integral becomes, setting $y(t) = q'(t) - q'_{cl}$ and noting $y(t)$ now has the boundary conditions $y(0) = y(T) = 0$,

$$G(T) = \int dq_0 e^{-\frac{i}{\hbar} V(x_0) - \frac{i}{\hbar} 2\omega(q_0 - q_{cl})^2 \frac{\sin^2 \omega T / 2}{\sin \omega T}} \int \mathcal{D}[y(t)] e^{\frac{i}{2\hbar} \int_0^T dt y(-\frac{\partial^2}{\partial t^2} - \omega^2)y}$$

Solving for the determinant of the remaining action $-\frac{1}{2} \int_0^T dt y(\frac{\partial^2}{\partial t^2} + \omega^2)y$ we solve the relevant Jacobi equation⁵⁵ $(\frac{\partial^2}{\partial t^2} + \omega^2)y = 0$ with initial conditions

$y(0) = 0$ and $y'(0) = 1$. This gives the prefactor

$$\sqrt{\frac{\omega}{2\pi i \hbar \sin \omega T}} \quad (\text{B.14})$$

Evaluating the q_0 integral, the final result is

$$\begin{aligned} G(T) &= \frac{1}{2i \sin \omega T/2} \\ &= e^{-i\omega T/2} \frac{1}{1 - e^{-i\omega T}} \\ &= \sum_{n=0}^{\infty} e^{-i(n+\frac{1}{2})\omega T - iTV(x_0)} \end{aligned} \quad (\text{B.15})$$

giving the excitation spectrum $E_n = \hbar\omega(n + \frac{1}{2})$ as expected.

Semiclassical approximation in field theory

This follows almost identically to the single particle case, with just a few technical points needing clarification. Suppose we have a field theory in 1+1 dimensions with the Lagrangian density

$$\mathcal{L}(x, t) = \frac{1}{2}(\partial_\mu \phi)^2 - U[\phi] \quad (\text{B.16})$$

Assume $\phi_{cl}(x)$ is a stationary extremum of this system. Expanding the action about this solution, $\phi \rightarrow \phi' + \phi_{cl}$

$$\mathcal{S} = S_{cl} + \frac{1}{2} \int dx \int dt (\partial_\mu \phi')^2 - \frac{\partial^2 U(\phi_{cl})}{\partial \phi^2} \phi'^2 \quad (\text{B.17})$$

Next, we integrate by parts to replace $(\partial_\mu \phi')^2 \rightarrow \phi'(-\frac{\partial^2}{\partial t^2} + \frac{\partial^2}{\partial x^2})\phi'$.

Assuming now that $\phi'(x, t)$ is separable, i.e. $\phi'(x, t) = f(x)g(t)$, we solve for the eigenvalues of the spatial portion

$$\left(-\frac{\partial^2}{\partial x^2} + \frac{\partial^2 U(\phi_{cl})}{\partial \phi^2}\right) f_r(x) = \omega_r^2 f_r(x) \quad (\text{B.18})$$

Assume that the $f_r(x)$ eigenfunctions form an orthonormal basis. Expressing the general solution $\phi'(x, t) = \sum_r f_r(x)g_r(t)$ so that the integration measure becomes $\prod_r \mathcal{D}[g_r(t)]$, the action becomes

$$\begin{aligned}
\mathcal{S} &= \frac{1}{2} \int dx \int dt \sum_r f_r(x) g_r(t) \sum_{r'} \left(-\frac{\partial^2}{\partial t^2} - \omega_{r'}^2 \right) f_{r'}(x) g_{r'}(t) \\
&= \sum_r \frac{1}{2} \int dt g_r(t) \left(-\frac{\partial^2}{\partial t^2} - \omega_r^2 \right) g_r(t)
\end{aligned} \tag{B.19}$$

by the orthonormality of the $f_r(x)$. Thus, the problem has separated into a product on r of equivalent single particle problems

$$G(T) = e^{\frac{i}{\hbar} S_{cl}} \prod_r \left(\int \mathcal{D}[g_r(t)] e^{\frac{i}{2\hbar} \int dt g_r(t) \left(-\frac{\partial^2}{\partial t^2} - \omega_r^2 \right) g_r(t)} \right) \tag{B.20}$$

which we know how to solve from the previous section.

The only remaining manipulation required is to note that

$$\prod_r \sum_{n_r} e^{-iT\omega_r(n_r + \frac{1}{2})} = \sum_{\{n_r\}} e^{-\frac{iT}{\hbar} \sum_r \hbar \omega_r(n_r + \frac{1}{2})} \tag{B.21}$$

where $\{n_r\}$ denotes a set of integers n_r .

B.1.2 Collective coordinates

Suppose the soliton exists in a system with translational symmetry. The soliton itself is a localized entity, and hence breaks this symmetry. The soliton must choose arbitrarily what coordinate to center on. This is an example of spontaneously broken symmetry.

This symmetry introduces to the quasiparticle spectrum a zero frequency mode associated with the soliton. While to first order presenting no problems, should we continue in the perturbative expansion, the energy denominators would develop artificial singularities.

In perturbing about the soliton solution, rather than as done previously via

$$\begin{aligned}
\phi &= \phi_0 + \sum_{n=0}^{\infty} a_n(t) \psi_n(x) \\
&= \phi_0 + a_0(t) \frac{d\phi_0}{dx} + \sum_{n=1}^{\infty} a_n(t) \psi_n(x)
\end{aligned} \tag{B.22}$$

where the $n = 0$ mode is the translation mode, rewrite the expansion as

$$\phi = \phi_0(x - X(t)) + \sum_{n=1}^{\infty} a_n(t) \psi_n(x) \tag{B.23}$$

where $X(t)$ is the collective coordinate associated to the translational invariance. This is completely equivalent if we expand $\phi_0(x - X(t))$ to first order in $X(t)$ and identify $a_0(t)$ with $-X(t)$.

Rewriting the Lagrangian in terms of this expansion, the potential terms, being translationally invariant by assumption, does not depend on $X(t)$. The kinetic term depends only on $\dot{X}(t)$.

We can introduce conjugate momenta to $X(t)$ and to the $a_n(t)$, denote these P and π_n , and transform to the classical Hamiltonian

$$H = P\dot{X}(t) + \sum_{n=1}^{\infty} \pi_n \dot{a}_n(t) - L \quad (\text{B.24})$$

Quantizing the soliton now follows exactly as quantization of a regular particle: we impose commutation relations on the various degrees of freedom

$$\begin{aligned} [X, P] &= i\hbar \\ [a_n, \pi_n] &= i\hbar \end{aligned} \quad (\text{B.25})$$

The quantized quasiparticles have a zero-point energy shifted by $\sum \frac{1}{2}\hbar\delta\omega_n$ that is attributed instead to the quantized soliton. That is, if the vacuum quasiparticle zero-point energy is $\sum \frac{1}{2}\hbar\omega_n$, while in the presence of a soliton becomes $\sum \frac{1}{2}\hbar(\omega_n + \delta\omega_n)$, the soliton is said to have the zero-point energy $\sum \frac{1}{2}\hbar\delta\omega_n$ while the quasiparticles are considered unchanged⁵⁰.

Appendix C

Spin path integrals

Consider a spin system $\Omega(t)$ with Hamiltonian H . The propagator for this spin to evolve from state Ω_i at time $t = 0$ to Ω_f at time $t = T$ is

$$K(\Omega_f, T; \Omega_i, 0) = \langle \Omega_f | \exp -\frac{i}{\hbar} H T | \Omega_i \rangle \quad (\text{C.1})$$

Inserting $N - 1$ resolutions of the identity³

$$\frac{2s+1}{4\pi} \int d\Omega |\Omega\rangle \langle \Omega| = 1 \quad (\text{C.2})$$

where a lower case s denotes the dimensionless spin (whereas, $S = \hbar s$), gives

$$K(\Omega_f, T; \Omega_i, 0) = \prod_{k=1}^{N-1} \left(\frac{2s+1}{4\pi} \int d\Omega_k \right) \langle \Omega_N | \exp -\frac{i}{\hbar} H \frac{T}{N} | \Omega_{N-1} \rangle \\ \langle \Omega_{N-1} | \cdots | \Omega_1 \rangle \langle \Omega_1 | \exp -\frac{i}{\hbar} H \frac{T}{N} | \Omega_0 \rangle$$

where $k = 0$ denotes the initial state and $k = N$ the final state. Define $\epsilon = T/N$.

Expand the exponential

$$\langle \Omega_{k+1} | \exp -\frac{i}{\hbar} H \frac{T}{N} | \Omega_k \rangle = \langle \Omega_{k+1} | \Omega_k \rangle \left(1 - \frac{i}{\hbar} \epsilon \frac{\langle \Omega_{k+1} | H | \Omega_k \rangle}{\langle \Omega_{k+1} | \Omega_k \rangle} + \mathcal{O}(\epsilon^2) \right)$$

Keeping terms to linear order in ϵ , the H term can be approximated at equal times: define $H(t_k) = \langle \Omega_{k+1} | H | \Omega_k \rangle$. Re-exponentiate the bracketed term to $\exp -\frac{i}{\hbar} \epsilon H(t_k)$.

The overlap of two coherent states, Ω_k and Ω_{k+1} is³

$$\langle \Omega_{k+1} | \Omega_k \rangle = \left(\frac{1 + \hat{\Omega}_{k+1} \cdot \hat{\Omega}_k}{2} \right)^s e^{-is\psi} \quad (\text{C.3})$$

where

$$\psi = 2 \tan^{-1} \left(\tan \left(\frac{\phi_{k+1} - \phi_k}{2} \right) \frac{\cos \frac{1}{2}(\theta_{k+1} + \theta_k)}{\cos \frac{1}{2}(\theta_{k+1} - \theta_k)} \right) + \xi_{k+1} - \xi_k \quad (\text{C.4})$$

and where ξ is a gauge dependent phase that we can ignore. The pre-factor is 1 to leading order and the phase can be approximated such that

$$\langle \Omega_{k+1} | \Omega_k \rangle = \exp \left(-is\epsilon \frac{\phi_{k+1} - \phi_k}{2} \cos \theta_k \right) \quad (\text{C.5})$$

All together, letting $N \rightarrow \infty$, we find the spin path integral

$$K(\Omega_f, T; \Omega_i, 0) = \int \mathcal{D}[\Omega(t)] \exp \left(-is \int_0^T dt \dot{\phi}(t) \cos \theta(t) - H(t) \right) \quad (\text{C.6})$$

Note that there are no spurious boundary terms as there are in the stereographic representation using z and z^* , as found, for example, by Solari⁶⁰.

C.1 The semiclassical approximation

Evaluation of spin path integral is non-trivial as evidenced by the series of papers suggesting various corrections. Klauder³² discussed the spin path integral in terms of conjugate variables ϕ and $S \cos \theta$ and first addressed the semiclassical approximation applied to the spin path integral. He claimed that to evaluate properly the trace of the propagator obtaining the excitation spectrum, real valued periodic orbits are required. However, simple counting of degrees of freedom, given two equations of motion (one for ϕ and another for $S \cos \theta$) with two initial and two final conditions, results in an overdetermined system. In fact, we are also trying to simultaneously specify both x and p at each boundary, disallowed by the familiar uncertainty principle.

Kuratsuji and Mizobuchi³⁵ note this overdeterminacy and claim only one of $\{x_i, x_f\}$ or $\{p_i, p_f\}$ needs specifying, the other being fixed by the equations of motion.

Solari⁶⁰ finds an additional pre-factor

$$\exp \frac{i}{2} \int_0^T dt \tilde{A}(t) \quad (\text{C.7})$$

where $\tilde{A}(t)$ is a time-dependent operator appearing in the action $z \tilde{A}(t) z^*$ where z is the spin coherent state in the stereographic projection. We won't worry about this correction since in our treatment there is no such term in the action.

Various authors⁵⁷ have even claimed that the spin path integral can only be properly evaluated discretely. However, a continuous version is reliable with the proper additional phase of Solari, as argued by Stone et. al.⁶³.

Below, we review the usage of the Jacobi equation for evaluating the path integral of a regular particle, then generalizing to the path integral over a field. Finally, we derive the analogous Jacobi equation for a spin path integral, following closely the work of Kuratsuji³⁵.

C.1.1 Coherent state path integral

In the classical limit, a spin coherent state $|\Omega(t)\rangle$ can be interpreted simply as a spin lying along the direction $\Omega(t)$.

The spin path integral, including the trace over periodic orbits, can be written as[†]

$$G(T) = \int_{\Omega(0)=\Omega(T)} \mathcal{D}[\Omega(t)] e^{\frac{i}{\hbar} \int_0^T dt - S\dot{\phi} \cos \theta - H[\Omega(t)]} \quad (\text{C.8})$$

This step is analogous to the quantum perturbations about a soliton solution. We now are solving for the quantum propagator for these perturbations.

Let $\phi_{cl}(t)$ and $\theta_{cl}(t)$ be a classical solution of this action (analogous to the simple harmonic oscillator solutions of the single particle case). Attempting to impose periodic boundary conditions results, in general, in an over-determined system of equations. Instead, we set only $\phi_{cl}(0) = \phi_{cl}(T)$ allowing the equations of motion to fix boundary conditions for $\theta_{cl}(t)$.

Expanding $\phi = \phi_{cl}(t) + x(t)$ and $S\theta = S\theta_{cl}(t) + y(t)$, the action becomes to second order variations (neglecting higher orders in keeping with the semiclassical approximation)

$$\mathcal{S} = \mathcal{S}_{cl} - \int dt \left(\dot{x}y + \frac{1}{2} (A(t)x^2 + 2B(t)xy + C(t)y^2) \right) \quad (\text{C.9})$$

where $A(t) = \frac{\partial^2 H}{\partial \phi^2}$, $B(t) = \frac{\partial^2 H}{\partial \phi \partial S \cos \theta}$ and $C(t) = \frac{\partial^2 H}{\partial (S \cos \theta)^2}$.

In the discrete version[‡], introducing the small timestep ϵ , we complete the square in y_k to obtain

[†]For the moment considering a single spin – the generalization to a field of spins follows identically to the treatment in Appendix B.

[‡]In arriving at this expression, note that in the discrete version there is actually an average of $y(t) \rightarrow \frac{y_k + y_{k-1}}{2}$ which under careful analysis gives boundary terms as found by Solari⁶⁰. We neglect these terms and approximate $\frac{y_k + y_{k-1}}{2} \approx y_k$.

$$\begin{aligned} \mathcal{S} = \mathcal{S}_{cl} - \frac{1}{2} \sum_{k=1}^N \epsilon C_k \left(y_k + \frac{B_k}{C_k} x_k y_k + \frac{(x_k - x_{k-1})}{\epsilon C_k} \right)^2 \\ + \epsilon x_k^2 \left(A_k - \frac{B_k^2}{C_k} + \frac{d}{dt} \left(\frac{B}{C} \right)_k \right) - \frac{(x_k - x_{k-1})^2}{\epsilon C_k} \end{aligned} \quad (\text{C.10})$$

Notice we use the extra integration over the periodic orbit coordinate θ_N to integrate over all N y_k 's; whereas, we only use $N-1$ integrations over the x_k 's. Impose the boundary conditions $x_0 = x_N = 0$.

For the general case, where we do not have the additional integration over boundary conditions, we must introduce this additional integration as an averaging over the final coordinate. This doesn't change the physics since this final coordinate is necessarily fixed by the equations of motion anyway.

The N Gaussian integrals over y_k give the pre-factors $\prod_{k=1}^N \sqrt{\frac{2\pi}{i\epsilon C_k}}$. The complete expression becomes

$$\begin{aligned} G(T) = \lim_{N \rightarrow \infty} \left(\frac{1}{2\pi} \right)^N \int d\phi_0 \left(\prod_{k=1}^{N-1} dx_k \right) \left(\prod_{k=1}^N \sqrt{\frac{2\pi}{\epsilon C_k}} \right) \\ \exp \frac{i}{2} \sum_{k=1}^N N \frac{x_k^2 - 2x_k x_{k-1} + x_{k-1}^2}{\epsilon C_k} - \epsilon x_k^2 a_k \end{aligned} \quad (\text{C.11})$$

where $a_k = A_k - \frac{B_k^2}{C_k} + \frac{d}{dt} \left(\frac{B}{C} \right)_k$.

The problem becomes that of solving for the determinant of the $(N-1) \times (N-1)$ matrix

$$-i \begin{pmatrix} \tilde{a}_1 & -\frac{1}{\epsilon C_2} & & & \\ -\frac{1}{\epsilon C_2} & \tilde{a}_2 & & & \\ & & \ddots & & \\ & & & \tilde{a}_{N-2} & -\frac{1}{\epsilon C_{N-1}} \\ & & & -\frac{1}{\epsilon C_{N-1}} & \tilde{a}_{N-1} \end{pmatrix}$$

where $\tilde{a}_k = -\epsilon a_k + \frac{1}{\epsilon C_k} + \frac{1}{\epsilon C_{k+1}}$.

Re-express the product of pre-factors from the y_k Gaussian integrals as

$$\prod_{k=1}^N \sqrt{\frac{2\pi}{i\epsilon C_k}} = \left(\det \begin{pmatrix} iC_2\epsilon & & & \\ & iC_3\epsilon & & \\ & & \ddots & \\ & & & iC_N\epsilon \end{pmatrix} i\epsilon C_1 \right)^{1/2}$$

and noting that $\det(AB) = \det(A)\det(B)$, we multiply the two matrices to yield

$$iC_1\epsilon \det \begin{pmatrix} C_2\epsilon\tilde{a}_1 & -1 & & \\ -\frac{C_3}{C_2} & C_3\epsilon\tilde{a}_2 & & \\ & & \ddots & \\ & & & C_{N-1}\epsilon\tilde{a}_{N-2} & -1 \\ & & & -\frac{C_N}{C_{N-1}} & C_N\epsilon\tilde{a}_{N-1} \end{pmatrix}$$

Denote the determinant of the submatrix ending in the k 'th row and column by D_k . We can then write down the recursion relation

$$\begin{aligned} D_k &= \epsilon C_{k+1}\tilde{a}_k D_{k-1} - \frac{C_{k+1}}{C_k} D_{k-2} \\ &= \left(1 + \frac{C_{k+1}}{C_k} - \epsilon^2 C_{k+1} \left(A_k - \frac{B_k^2}{C_k} + \frac{d}{dt} \left(\frac{B}{C}\right)_k\right)\right) D_{k-1} - \frac{C_{k+1}}{C_k} D_{k-2} \end{aligned}$$

Letting D_k be a function of $k\epsilon$, this can be rewritten

$$\begin{aligned} \frac{D_k - 2D_{k-1} + D_{k-2}}{\epsilon^2} &= \frac{(C_{k+1} - C_k)(D_{k-1} - D_{k-2})}{C_k \epsilon^2} \\ &\quad - C_{k+1} \left(A_k - \frac{B_k^2}{C_k} + \frac{d}{dt} \left(\frac{B}{C}\right)_k\right) D_{k-1} \end{aligned}$$

or in a continuum limit

$$\frac{d^2 D}{dt^2} = \frac{1}{C} \frac{dC}{dt} \frac{dD}{dt} - CD \left(A - \frac{B^2}{C} + \frac{d}{dt} \left(\frac{B}{C}\right)\right) \quad (\text{C.12})$$

The initial conditions on D can be found directly from the first and second submatrix determinants

$$\begin{aligned} D_1 &= iC_1\epsilon C_2\epsilon\tilde{a}_1 \\ \frac{D_2 - D_1}{\epsilon} &= iC_1 \left(C_3\epsilon\tilde{a}_2 C_2\epsilon\tilde{a}_1 - \frac{C_3}{C_2} - C_2\epsilon\tilde{a}_1\right) \end{aligned}$$

giving, in the limit of $\epsilon \rightarrow 0$, $D(0) = 0$ and $\dot{D}(0) = iC(0)$. But this is equivalent to the system of equations from the original formulation

$$\begin{aligned} \frac{dy}{dt} &= Ax + By \\ \frac{dx}{dt} &= -Bx - Cy \end{aligned} \quad (\text{C.13})$$

with initial conditions $x(0) = 0$ and $y(0) = -1$ after eliminating $y(t)$ and setting $ix(t) = D(t)$. Thus the required determinant is $ix(T)$.

C.1.2 Spectrum of a ferromagnetic plane of spins

The action of a ferromagnetic plane of spins with easy plane anisotropy in a continuum limit is

$$\mathcal{S} = S \int \frac{d^2r}{a^2} \int dt \dot{\phi} \theta - \frac{c}{2} \left(-\phi \nabla^2 \phi - \theta \nabla^2 \theta + \frac{\theta^2}{r_v^2} \right) \quad (\text{C.14})$$

where the various constants are as defined in Chapter 2.

Choose a set of spatial eigenfunctions such that $\nabla^2 \rightarrow -k^2$ and $\int \frac{d^2r}{a^2} f_{k'} f_k = \delta^2(k' - k)$.

Thus, the integration measure becomes a product over k states, now decoupled, leaving within the time integral of the action (note S was factored out into the integration measure)

$$\frac{1}{2} \begin{pmatrix} \phi & \theta \end{pmatrix} \begin{pmatrix} -c\nabla^2 & -\frac{\partial}{\partial t} \\ \frac{\partial}{\partial t} & -c \left(\nabla^2 - \frac{1}{r_v^2} \right) \end{pmatrix} \begin{pmatrix} \phi \\ \theta \end{pmatrix} \quad (\text{C.15})$$

where $\omega = ckQ$. The periodic classical solutions can be written

$$\begin{pmatrix} \phi_k(t) \\ \theta_k(t) \end{pmatrix} = A \begin{pmatrix} \cos \omega_k t \\ -\frac{k}{Q} \sin \omega_k t \end{pmatrix} + B \begin{pmatrix} \sin \omega_k t \\ \frac{k}{Q} \cos \omega_k t \end{pmatrix} \quad (\text{C.16})$$

with the periodicity condition on $\phi_k(t)$ imposing identical conditions on A and B as in (B.12), with $q_0 - q_{cl} \rightarrow \phi_{k0}$. Note that the periodicity condition was previously $\phi(x, 0) = \phi(x, T) = \phi_0$; however, after the transformation to diagonalize the equations in k , each coefficient ϕ_{k0} must now be periodic and integrated over.

The classical action for these periodic orbits becomes

$$\mathcal{S}_{cl} = -2 \frac{k}{Q} \phi_{k0}^2 \frac{\sin^2 \omega_k T / 2}{\sin \omega_k T} \quad (\text{C.17})$$

The perturbed action has the same form as the original linearized action above, (C.14). Calling the small perturbations in ϕ , x , and those in θ , y , we need a solution such that $x(0) = 0$ and $y(0) = 1$ (the change of sign here arises from linearizing $\cos \theta \rightarrow -\theta$ in the Berry phase term). This corresponds to

$$\begin{pmatrix} x(t) \\ y(t) \end{pmatrix} = \begin{pmatrix} \frac{Q}{k} \sin \omega_k t \\ \cos \omega_k t \end{pmatrix} \quad (\text{C.18})$$

and the determinant evaluates to $ix(T) = i\frac{Q}{k} \sin \omega_k T$.

Bibliography

1. Y. Aharonov and D. Bohm. Significance of electromagnetic potentials in the quantum theory. *Phys. Rev.*, 115, 485 (1959).
2. P. Ao and D. J. Thouless. Berry's phase and the Magnus force for a vortex line in a superconductor. *Phys. Rev. Lett.*, 70, 2158 (1993).
3. Assa Auerbach. *Interacting electrons and quantum magnetism*. Springer-Verlag, New York (1994).
4. M. V. Berry. Quantal phase factors accompanying adiabatic changes. *Proc. R. Soc. Lond. A*, 392, 45 (1984).
5. A. O. Caldeira and A. J. Leggett. Path integral approach to quantum brownian motion. *Physica A*, 121, 587 (1983).
6. S. Coleman. Classical lumps and their quantum descendants. *Phys. Rep. C*, 21, 227 (1975).
7. J. F. Currie, J. A. Krumhansl, A. R. Bishop, and S. E. Trullinger. Statistical mechanics of one-dimensional solitary-wave-bearing scalar fields: Exact results and ideal-gas phenomenology. *Phys. Rev. B*, 22, 477 (1980).
8. R. F. Dashen, B. Hasslacher, and A. Neveu. Nonperturbative methods and extended-hadron models in field theory. I. Semiclassical functional methods. *Phys. Rev. D*, 10, 4114 (1974).
9. E. Demircan, P. Ao, and Q. Niu. Interactions of collective excitations with vortices in superfluid systems. *Phys. Rev. B*, 52, 476 (1995).
10. P.A.M. Dirac. The fundamental equations of quantum mechanics. *Proc. Roy. Soc. A*, 109, 642 (1925).
11. J. M. Duan and A. J. Leggett. Inertial mass of a moving singularity in a fermi superfluid. *Phys. Rev. Lett.*, 68, 1216 (1992).
12. M. Dube and P. C. E. Stamp. Effects if phonons and nuclear spins on the tunneling of a domain wall. *J. Low Temp. Phys.*, 110(3), 779 (1998).
13. A. L. Fetter. Scattering of Sound by a Classical Vortex. *Phys. Rev.*, 136, A1488 (1964).

-
14. R. P. Feynman. Application of quantum mechanics to liquid helium. In C. J. Gorter, editor, *Progress in Low Temperature Physics*, volume 1. North-Holland, Amsterdam (1955).
 15. R. P. Feynman. *Statistical mechanics : a set of lectures*. Westview Press, Boulder (1998, c1972).
 16. R. P. Feynman and A. R. Hibbs. *Quantum mechanics and path integrals*. McGraw-Hill, New York (1965).
 17. R. P. Feynman and F. L. Vernon. The theory of a general quantum system interacting with a linear dissipative system. *Annals of Physics*, 24, 118 (1963).
 18. J.-L. Gervais and B. Sakita. Extended particles in quantum field theories. *Phys. Rev. D*, 11, 2943 (1975).
 19. J. Goldstone. *Nuovo Cimento*, 19, 154 (1961).
 20. J. Goldstone and R. Jackiw. Quantization of nonlinear waves. *Phys. Rev. D*, 11, 1486 (1975).
 21. M. E. Gouvea, G. M. Wysin, A. R. Bishop, and F. G. Mertens. Vortices in the classical two-dimensional anisotropic Heisenberg model. *Phys. Rev. B*, 39, 11840 (1989).
 22. H. E. Hall and J. R. Hook. Comment on ‘Magnus and Iordanskii forces in superfluids’. *Phys. Rev. Lett.*, 80, 4356 (1998).
 23. H. E. Hall and W. F. Viven. The rotation of liquid helium II. I. Experiments on the propagation of second sound in uniformly rotating helium II. *Proc. R. Soc. London, Ser. A*, 238, 204 (1956).
 24. H. E. Hall and W. F. Viven. The rotation of liquid helium II. II. The theory of mutual friction in uniformly rotating helium II. *Proc. R. Soc. London, Ser. A*, 238, 215 (1956).
 25. D. L. Huber. Dynamics of spin vortices in two-dimensional planar magnets. *Phys. Rev. B*, 26, 3758 (1982).
 26. S. V. Iordanskii. On the mutual friction between the normal and superfluid components in a rotating Bose gas. *Ann. Phys.*, 29, 335 (1964).
 27. S. V. Iordanskii. Mutual friction force in a rotating bose gas. *Sov. Phys. JETP*, 22, 160 (1966).
 28. B. A. Ivanov, H. J. Schnitzer, F. G. Mertens, and G. M. Wysin. Magnon modes and magnon-vortex scattering in two-dimensional easy-plane ferromagnets. *Phys. Rev. B*, 58, 8464 (1998).

-
29. B. A. Ivanov and I. A. Yastremsky. Scattering of a radially symmetric spin wave on a magnetic vortex in a two-dimensional easy-plane ferromagnet. *Low. Temp. Phys.*, 26(5), 341 (2000).
 30. R. Jackiw and C. Rebbi. Solitons with fermion number $1/2$. *Phys. Rev. D*, 13, 3398 (1976).
 31. John David Jackson. *Classical electrodynamics*. John Wiley & Sons, New York, third edition (1998, c1999).
 32. J. R. Klauder. Path integrals and stationary-phase approximations. *Phys. Rev. D*, 19, 2349 (1979).
 33. J. M. Kosterlitz and D. J. Thouless. Ordering, metastability and phase transitions in two-dimensional systems. *J. Phys. C: Solid State Phys.*, 6, 1181 (1973).
 34. J. A. Krumhansl and J. R. Schrieffer. Dynamics and statistical mechanics of a one-dimensional model Hamiltonian for structural phase transitions. *Phys. Rev. B*, 11, 3535 (1975).
 35. H. Kuratsuji and Y. Mizobuchi. A semiclassical treatment of path integrals for the spin system. *J. Math. Phys.*, 22, 757 (1981).
 36. Fluid Dynamics Laboratory. Online. <http://www.fluid.tue.nl/WDY/vort/> (July 24, 2004).
 37. L. D. Landau and E. M. Lifshitz. *The classical theory of fields*, volume 2 of *Course of Theoretical Physics*. Pergamon Press, Oxford; New York, third edition (1971).
 38. L. D. Landau and E. M. Lifshitz. *Quantum mechanics: Non-relativistic theory*, volume 3 of *Course of Theoretical Physics*. Pergamon Press, Oxford; New York, third edition (1989, c1977).
 39. F. G. Mertens and A. R. Bishop. Dynamics of vortices in two-dimensional magnets. *cont-mat/9903037* (1999).
 40. F. G. Mertens, A. R. Bishop, G. M. Wysin, and C. Kawabata. Vortex signatures in dynamic structure factors for two-dimensional easy-plane ferromagnets. *Phys. Rev. Lett.*, 59, 117 (1987).
 41. F. G. Mertens, H. J. Schnitzer, and A. R. Bishop. Hierarchy of equations of motion for nonlinear coherent excitations applied to magnetic vortices. *Phys. Rev. B*, 56, 2510 (1997).
 42. A. H. Castro Neto and A. O. Caldeira. Transport properties of solitons. *Phys. Rev. E*, 48, 4037 (1993).
 43. A. V. Nikiporov and E. B. Sonin. Dynamics of magnetic vortices in a planar ferromagnet. *Sov. Phys. JETP*, 58, 373 (1983).

-
44. V. Novosad, M. Grimsditch, K. Yu. Guslienko, P. Vavassori, Y. Otani, and S. D. Bader. Spin excitations of magnetic vortices in ferromagnetic nanodots. *Phys. Rev. B*, 66, 52407 (2002).
 45. L. Onsager. *Nuovo Cimento Suppl.*, 6, 249 (1949).
 46. R. Peierls. *Surprises in Theoretical Physics*. Princeton Series in Physics, Princeton (1979).
 47. R. F. Peierls. *Quantum theory of solids*. Clarendon Press, Oxford (1955).
 48. A. R. Pereira, F. O. Coelho, and A. S. T. Pires. Normal modes of vortices in easy-plane antiferromagnets: Exact results and Born approximation. *Phys. Rev. B*, 54, 6084 (1996).
 49. R. Rajaraman. *Phys. Rep. C*, 21, 227 (1975).
 50. R. Rajaraman. *Solitons and instantons, An introduction to solitons and instantons in quantum field theory*. North Holland Publishing, Amsterdam; New York; Oxford (1982).
 51. J. P. Rodriguez. Quantized topological point defects in two-dimensional antiferromagnets. *Phys. Rev. B*, 39, 2906 (1989).
 52. C. A. Ross. Patterned magnetic recording media. *Annu. Rev. Mater. Res.*, 31, 203 (2001).
 53. John Scott Russell. Report on waves. In *Report of the fourteenth meeting of the British Association for the Advancement of Science*. London (1845).
 54. B. Sakita. *Quantum theory of many-variable systems and fields*. World Scientific, Singapore (1985).
 55. Lawrence S. Schulman. *Techniques and applications of path integrals*. Wiley, New York (c1981).
 56. D. D. Sheka, I. A. Yastremsky, B. A. Ivanov, G. M. Wysin, and F. G. Mertens. Amplitudes for magnon scattering by vortices in two-dimensional weakly easy-plane. *cond-mat/0309640* (2003).
 57. Junya Shibata and Shin Takagi. Method of collective degrees of freedom in spin coherent state path integral. *cond-mat/0105365* (2001).
 58. T. Shinjo, T. Okuno, R. Hassdorf, K. Shigeto, and T. Ono. Magnetic vortex core observation in circular dots of permalloy. *Science*, 289, 930 (2000).
 59. J. C. Slonczewski. Motions of magnetic vortex solitons. In J. Rauluszkievicz, H. Szymczak, and H. K. Lachowicz, editors, *Physics of Magnetic Materials*. World Scientific, Singapore (1985).
 60. H. D. Solari. Semiclassical treatment of spin system by means of coherent state. *J. Math. Phys.*, 28, 1097 (1987).

-
61. E. B. Sonin. Magnus force in superfluids and superconductors. *Phys. Rev. B*, 55, 485 (1997).
 62. P. C. E. Stamp. Quantum dynamics and tunneling of domain walls in ferromagnetic insulators. *Phys. Rev. Lett.*, 66, 2802 (1991).
 63. Michael Stone, Kee-Su Park, and Anupam Garg. The semiclassical propagator for spin coherent states. *J. Math. Phys.*, 41, 8025 (2000).
 64. W. P. Su, J. R. Schrieffer, and A. J. Heeger. Solitons in polyacetylene. *Phys. Rev. Lett.*, 42, 1698 (1979).
 65. A. A. Thiele. Steady-state motion of magnetic domains. *Phys. Rev. Lett.*, 30, 230 (1973).
 66. A. A. Thiele. Applications of the gyrocoupling vector and dissipation dyadic in the dynamics of magnetic domains. *J. Appl. Phys.*, 45(1), 377 (1974).
 67. D. J. Thouless, P. Ao, and Q. Niu. Transverse force on a quantized vortex in a superfluid. *Phys. Rev. Lett.*, 76, 3758 (1996).
 68. E. Tomboulis. Canonical quantization of nonlinear waves. *Phys. Rev. D*, 12, 1678 (1975).
 69. A. R. Volkel, F. G. Mertens, G. M. Wysin, A. R. Bishop, and H. J. Schnitzer. Collective variable approach for a magnetic n-vortex system. In K. H. Spatscheck and F. G. Mertens, editors, *Nonlinear Coherent Structures in Physics and Biology*. Plenum Press, New York (1994).
 70. G. E. Volovik. Comment on ‘Transverse force on a quantized vortex in a superfluid’. *Phys. Rev. Lett.*, 77, 4687 (1996).